

Rabbit anti-Human AIF Polyclonal Antibody | anti-AIF antibody

AIF Antibody

CATALOG # AAA306683	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunocytochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

AIF; Polyclonal Antibody; AIF Antibody; anti-AIF antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Ion exchange chromatography purified
Form/Format	Supplied in PBS containing 0.02% sodium azide.
Sequence Length	261
Applicable Applications for anti-AIF antibody	ICC (Immunocytochemistry), WB (Western Blot), ELISA
Immunogen Type	Peptide
Immunogen Description	Raised against a peptide corresponding to amino acids near the amino terminus of mature human AIF.
Target Name	AIF
Preparation and Storage	Can be stored at -20 degree C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

IMAGES


ICC (Immunocytochemistry)

Immunocytochemistry of AIF in Jurkat cells with AIF antibody at 2 ug/mL.


WB (Western Blot)

Western blot analysis of AIF in K562 cell lysate with AIF antibody at 1 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for anti-AIF antibody	Apoptosis is characterized by several morphological nuclear changes including chromatin condensation and nuclear fragmentation. These changes are triggered by the activation of members of caspase family, caspase activated DNase, and several novel proteins. A novel gene, the product of which causes chromatin condensation and DNA fragmentation, was recently identified, cloned, and designated apoptosis inducing factor (AIF). Like the critical molecules, cytochrome c and caspase-9, in apoptosis, AIF localizes in mitochondria. AIF translocates to the nucleus when apoptosis is induced and induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. AIF induces chromatin condensation and DNA fragmentation, which are the hallmarks of apoptosis, of the isolated nucleus and the nucleus in live cells by microinjection. AIF is highly conserved between human and mouse and widely expressed.
Product Categories/Family for anti-AIF antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	195927004
NCBI GeneID	9131
NCBI Accession #	NP_001124318.1
NCBI GenBank Nucleotide #	NM_001130846.2
UniProt Accession #	O95831; Q1L6K4; Q1L6K6; Q2QKE4; Q5JUZ7; Q6I9X6; Q9Y3I3; Q9Y3I4; A4QPB4; B1ALN1; B2RB08; D3DTE9
Molecular Weight	26,033 Da
NCBI Official Full Name	apoptosis-inducing factor 1, mitochondrial isoform 4
NCBI Official Synonym Full Names	apoptosis-inducing factor, mitochondrion-associated, 1
NCBI Official Symbol	AIFM1
NCBI Official Synonym Symbols	AIF; CMT2D; CMTX4; COWCK; NADMR; NAMSD; PDCD8; COXPD6
NCBI Protein Information	apoptosis-inducing factor 1, mitochondrial
UniProt Protein Name	Apoptosis-inducing factor 1, mitochondrial
UniProt Gene Name	AIFM1
UniProt Synonym Gene Names	AIF; PDCD8
UniProt Entry Name	AIFM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/aif-antibody-immunocytochemistry-western-blot-elisa/306683>