

Rabbit anti-Human MCM7 Polyclonal Antibody | anti-MCM7 antibody

MCM7 antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA306691	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MCM7; Polyclonal Antibody; MCM7 antibody; anti-MCM7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Purified by antigen-affinity chromatography.
Form/Format	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Sequence Length	543
Applicable Applications for anti-MCM7 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 14 and 275 of MCM7
Target Name	MCM7
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using MCM7 antibody at 1: 200 dilution.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Cal27 Xenograft, using MCM7 antibody at 1: 100 dilution.



Application Data

Sample (30 ug of whole cell lysate) A: H1299 7.5% SDS PAGE Primary antibody diluted at 1:1000

ADDITIONAL INFORMATION

Related Product Information for anti-MCM7 antibody	The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumorsuppressor protein RB1/RB. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq]
Product Categories/Family for anti-MCM7 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469922
NCBI GeneID	4176
NCBI Accession #	NP_877577
NCBI GenBank Nucleotide #	NM_182776.2
Molecular Weight	60,643 Da
NCBI Official Full Name	DNA replication licensing factor MCM7 isoform 2
NCBI Official Synonym Full Names	minichromosome maintenance complex component 7
NCBI Official Symbol	MCM7
NCBI Official Synonym Symbols	MCM2; CDC47; P85MCM; P1CDC47; PNAS146; PPP1R104; P1.1-MCM3
NCBI Protein Information	DNA replication licensing factor MCM7
UniProt Protein Name	DNA replication licensing factor MCM7
UniProt Gene Name	MCM7
UniProt Synonym Gene Names	CDC47; MCM2
UniProt Entry Name	MCM7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

