

Rabbit HDAC5 Polyclonal Antibody | anti-HDAC5 antibody

HDAC5 (Phospho-Ser498) Antibody

CATALOG # AAA307132	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

HDAC5; Polyclonal Antibody; HDAC5 (Phospho-Ser498) Antibody; HD5; anti-HDAC5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of HDAC5 only when phosphorylated at serine498.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography use
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	1123
Applicable Applications for anti-HDAC5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 498 (T-Q-S(p)-S-P) derived from Human HDAC5/7.
Modification	Phospho-Ser498
Target Name	HDAC5
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using HDAC5(Phospho-Ser498) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from 293 cells untreated or treated with serum starvation using HDAC5(Phospho-Ser498) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-HDAC5 antibody	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by HDAC5 belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-HDAC5 antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	62750349
NCBI GeneID	10014
NCBI Accession #	NP_001015053.1
NCBI GenBank Nucleotide #	NM_001015053.1
UniProt Accession #	Q9UQL6; O60340; O60528; Q96DY4; C9JFV9
Molecular Weight	122,049 Da
NCBI Official Full Name	histone deacetylase 5 isoform 3
NCBI Official Synonym Full Names	histone deacetylase 5
NCBI Official Symbol	HDAC5
NCBI Official Synonym Symbols	HD5; NY-CO-9
NCBI Protein Information	histone deacetylase 5
UniProt Protein Name	Histone deacetylase 5
UniProt Gene Name	HDAC5
UniProt Synonym Gene Names	KIAA0600; HD5
UniProt Entry Name	HDAC5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hdac5-antibody-immunohistochemistry-western-blot-elisa/307132>