

Rabbit TLR7 Polyclonal Antibody | anti-TLR7 antibody

TLR7 antibody

CATALOG # AAA307292	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

TLR7; Polyclonal Antibody; TLR7 antibody; anti-TLR7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total TLR7 antibody.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	1049
Applicable Applications for anti-TLR7 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human TLR7.
Target Name	TLR7
Preparation and Storage	Store at -20 degree C

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of MCF7 cell using TLR7 antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of Ramos cell lysate using TLR7 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-TLR7 antibody	Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in Drosophila, play a pivotal role in innate immune responses (1-3). TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF-kB and subsequent regulation of immune and inflammatory genes. The TLRs and members of the IL-1 receptor family share a conserved stretch of approximately 200 amino acids known as the TIR domain. Upon activation, TLRs associate with a number of cytoplasmic adaptor proteins containing TIR domains including MyD88 (myeloid differentiation factor), MAL/TIRAP (MyD88-adaptor-like/TIR-associated protein), TRIF (Toll-receptor-associated activator of interferon), and TRAM (Toll-receptor-associated molecule). This association leads to the recruitment and activation of IRAK1 and IRAK4, which form a complex with TRAF6 to activate TAK1 and IKK. Activation of IKK leads to the degradation of IκB that normally maintains NF-kB inactivity by sequestering it in the cytoplasm.
Product Categories/Family for anti-TLR7 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	7706093
NCBI GeneID	51284
NCBI Accession #	NP_057646.1
NCBI GenBank Nucleotide #	NM_016562.3
UniProt Accession #	Q9NYK1; Q9NR98; D1CS69
Molecular Weight	120,922 Da
NCBI Official Full Name	toll-like receptor 7
NCBI Official Synonym Full Names	toll-like receptor 7
NCBI Official Symbol	TLR7
NCBI Official Synonym Symbols	TLR7-like
NCBI Protein Information	toll-like receptor 7
UniProt Protein Name	Toll-like receptor 7
UniProt Gene Name	TLR7
UniProt Entry Name	TLR7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tlr7-antibody-immunofluorescence-western-blot-elisa/307292>