

Rabbit anti-Human, Mouse MCM4 Polyclonal Antibody | anti-MCM4 antibody

MCM4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA307402	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MCM4; Polyclonal Antibody; MCM4 Antibody; CDC21; CDC54; hCdc21; P1-CDC21; anti-MCM4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total MCM4 protein.
Purity/Purification	Antigen affinity purification.
Form/Format	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Concentration	1.4 mg/ml (varies by lot)
Sequence Length	863
Applicable Applications for anti-MCM4 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide corresponding to a region derived from internal residues of human Minichromosome maintenance complex component 4
Target Name	MCM4
Preparation and Storage	Store at -20 degree C

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human brain tissue using at dilution 1/40.



Application Data

Gel: 8%SDS-PAGE Lysates (from left to right): NIH/3T3, HeLa and K562 cell Amount of lysate: 50ug per lane Primary antibody: 1/700 dilution Secondary antibody dilution: 1/8000 Exposure time: 10 seconds

ADDITIONAL INFORMATION

Related Product Information for anti-MCM4 antibody	The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 6 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of this protein by CDC2 kinase reduces the DNA helicase activity and chromatin binding of the MCM complex. This gene is mapped to a region on the chromosome 8 head-to-head next to the PRKDC/DNA-PK, a DNA-activated protein kinase involved in the repair of DNA double-strand breaks. Alternatively spliced transcript variants encoding the same protein have been reported.
Product Categories/Family for anti-MCM4 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469919
NCBI GeneID	4173
NCBI Accession #	NP_005905.2
NCBI GenBank Nucleotide #	NM_005914.3
UniProt Accession #	P33991; Q8NEH1; Q99658
Molecular Weight	96,558 Da
NCBI Official Full Name	DNA replication licensing factor MCM4
NCBI Official Synonym Full Names	minichromosome maintenance complex component 4
NCBI Official Symbol	MCM4
NCBI Official Synonym Symbols	NKCD; CDC21; CDC54; NKGCD; hCdc21; P1-CDC21
NCBI Protein Information	DNA replication licensing factor MCM4
UniProt Protein Name	DNA replication licensing factor MCM4
UniProt Gene Name	MCM4
UniProt Synonym Gene Names	CDC21
UniProt Entry Name	MCM4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mcm4-antibody-immunohistochemistry-western-blot-elisa/307402>