

Rabbit anti-Human FCGR2A Polyclonal Antibody | anti-FCGR2A antibody

FCGR2A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA307612	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

FCGR2A; Polyclonal Antibody; FCGR2A Antibody; CD32; CD32A; CDw32; FCG2; FCGR2; anti-FCGR2A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total FCGR2A protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	317
Applicable Applications for anti-FCGR2A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human FCGR2A.
Target Name	FCGR2A
Preparation and Storage	Store at -20 degree C

IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of HepG2 cell using FCGR2A antibody. Blue: DAPI for nuclear staining.


WB (Western Blot)

Western blot analysis of extracts of Jurkat cell lines, using FCGR2A antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-FCGR2A antibody	CD32 (also designated Fc gamma RII) is a low affinity receptor for the Fc fragment of aggregated IgG (1,2). CD32 is responsible for the clearance of immunocomplexes by macrophages and also plays an important role in the regulation of antibody production by B cells (1-4). IgG can noncooperatively bind either one or two highly glycosylated CD32 molecules, and this binding delivers.3 a negative signal for B cells (1,2,5). CD32 exists as several isoforms that are produced by alternative splicing of three distinct genes, A, B, and C (2,6). These isoforms are designated FcγRIIA, FcγRIIB1, FcγRIIB3, and FcγRIIC (1,2,6). All isoforms are present on monocytes, placental trophoblasts and endothelial cells (1,6). In addition, the FcγRIIB forms are present on B lymphocytes, and the FcγRIIA and FcγRIIC forms are found on neutrophils (1,6).
Product Categories/Family for anti-FCGR2A antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	210031822
NCBI GeneID	2212
NCBI Accession #	NP_001129691.1
NCBI GenBank Nucleotide #	NM_001136219.1
UniProt Accession #	P12318; Q8WUN1; Q8WW64
Molecular Weight	34,930 Da
NCBI Official Full Name	low affinity immunoglobulin gamma Fc region receptor II-a isoform 1
NCBI Official Synonym Full Names	Fc fragment of IgG, low affinity IIa, receptor (CD32)
NCBI Official Symbol	FCGR2A
NCBI Official Synonym Symbols	CD32; FCG2; FcGR; CD32A; CDw32; FCGR2; IGFR2; FCGR2A1
NCBI Protein Information	low affinity immunoglobulin gamma Fc region receptor II-a
UniProt Protein Name	Low affinity immunoglobulin gamma Fc region receptor II-a
UniProt Gene Name	FCGR2A
UniProt Synonym Gene Names	CD32; FCG2; FCGR2A1; IGFR2; IgG Fc receptor II-a; Fc-gamma-RIIa; FcRII-a
UniProt Entry Name	FCG2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fcgr2a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/307612>