

Rabbit 4E-BP1 Polyclonal Antibody | anti-4E-BP1 antibody

4E-BP1 (Ab-45) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA308246	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

4E-BP1; Polyclonal Antibody; 4E-BP1 (Ab-45) Antibody; EIF4EBP1; PHAS-1; anti-4E-BP1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total 4E-BP1 protein.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	118
Applicable Applications for anti-4E-BP1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.43~47 (S-T-T-P-G) derived from Human 4E-BP1.
Target Name	4E-BP1
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.


IMAGES


IF (Immunofluorescence)

Immunofluorescence staining of methanol-fixed HeLa cells using 4E-BP1(Ab-45) Antibody.


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using 4E-BP1(Ab-45) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from 293 and HepG2 cells using 4E-BP1(Ab-45) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-4E-BP1 antibody	4E-BP1 encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.
Product Categories/Family for anti-4E-BP1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758258
NCBI GeneID	1978
NCBI Accession #	NP_004086.1
NCBI GenBank Nucleotide #	NM_004095.3
UniProt Accession #	Q13541; Q6IBN3; B2R502; D3DSW8
Molecular Weight	12,580 Da
NCBI Official Full Name	eukaryotic translation initiation factor 4E-binding protein 1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 4E binding protein 1
NCBI Official Symbol	EIF4EBP1
NCBI Official Synonym Symbols	BP-1; 4EBP1; 4E-BP1; PHAS-I
NCBI Protein Information	eukaryotic translation initiation factor 4E-binding protein 1
UniProt Protein Name	Eukaryotic translation initiation factor 4E-binding protein 1
UniProt Gene Name	EIF4EBP1
UniProt Synonym Gene Names	4E-BP1; eIF4E-binding protein 1; PHAS-I
UniProt Entry Name	4EBP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/4e-bp1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/308246>