

Rabbit anti-Human Kininogen-1 Polyclonal Antibody | anti-KNG1 antibody

Kininogen-1 Polyclonal Antibody

CATALOG # AAA309120	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Kininogen-1; Polyclonal Antibody; Kininogen-1 Polyclonal Antibody; Alpha-2-thiol proteinase inhibitor; Fitzgerald factor; High molecular weight kininogen; Williams-Fitzgerald-Flaujeac factor; KNG1; BDK; KNG; anti-KNG1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total Kininogen-1 polyclonal antibody.
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	427
Applicable Applications for anti-KNG1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Human Kininogen-1 protein
Immunogen Type	Protein
Preparation and Storage	Store at -20 degree C

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded humankidney using at dilution of 1:50.



WB (Western Blot)

All lanes: Kininogen-1 antibody at 2ug/ml

ADDITIONAL INFORMATION

Related Product Information for anti-KNG1 antibody	Kininogens are inhibitors of thiol proteases; (2) HMW-kininogen plays an important role in blood coagulation by helping to position optimally prekallikrein and factor XI next to factor XII; (3) HMW-kininogen inhibits the thrombin- and plasmin-induced aggregation of thrombocytes; (4) the active peptide bradykinin that is released from HMW-kininogen shows a variety of physiological effects: (4A) influence in smooth muscle contraction, (4B) induction of hypotension, (4C) natriuresis and diuresis, (4D) decrease in blood glucose level, (4E) it is a mediator of inflammation and causes (4E1) increase in vascular permeability, (4E2) stimulation of nociceptors (4E3) release of other mediators of inflammation (e.g. prostaglandins), (4F) it has a cardioprotective effect (directly via bradykinin action, indirectly via endothelium-derived relaxing factor action); (5) LMW-kininogen inhibits the aggregation of thrombocytes; (6) LMW-kininogen is in contrast to HMW-kininogen not involved in blood clotting.
Product Categories/Family for anti-KNG1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504893
NCBI GeneID	3827
NCBI Accession #	NP_000884.1
NCBI GenBank Nucleotide #	NM_000893.3
UniProt Accession #	P01042; P01043; Q53EQ0; Q6PAU9; Q7M4P1; A8K474; B2RCR2; C9JEX1
Molecular Weight	43,822 Da
NCBI Official Full Name	kininogen-1 isoform 2
NCBI Official Synonym Full Names	kininogen 1
NCBI Official Symbol	KNG1
NCBI Official Synonym Symbols	BK; BDK; KNG
NCBI Protein Information	kininogen-1
UniProt Protein Name	Kininogen-1
UniProt Gene Name	KNG1
UniProt Synonym Gene Names	BDK; KNG; HMWK
UniProt Entry Name	KNG1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/kininogen-1-antibody-immunohistochemistry-western-blot-elisa/309120>