

Rabbit MITF Polyclonal Antibody | anti-MITF antibody

Phospho-MITF (Ser180/73) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30956	Rabbit	Human, Mouse, Rat, Monkey	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA


PRODUCT SYNONYMS

MITF; Polyclonal Antibody; Phospho-MITF (Ser180/73) Antibody; BHLHE32; bHLHe32; Class E basic helix-loop-helix protein 32; CMM8; Homolog of mouse microphthalmia; Mi; Microphthalmia associated transcription factor; Microphthalmia; mouse; homolog of; Microphthalmia-associated transcription factor; MITF_HUMAN; mitfa; nacre; WS2; WS2A; z3A.1; anti-MITF antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-MITF (Ser180) antibody detects endogenous levels of MITF only when phosphorylated at Serine 180
Purity/Purification	Peptide affinity purification
Form/Format	Supplied in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Sequence Length	419
Applicable Applications for anti-MITF antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 IF/ICC: 1:100-1:500 ELISA (peptide): 1:20000-1:40000 *The optimal dilution should be determined by the end user. IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.
Immunogen	A synthesized peptide derived from human MITF around the phosphorylation site of Ser180
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)

AAA30956 staining HepG2 cells(4h of LPS treatment) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AAA30956 1:200) and mouse anti-beta tubulin Ab(1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab. The nuclear counter stain is DAPI(blue).



IHC (Immunohistochemistry)

AAA30956 at 1/100 staining rat ovarian tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)

AAA30956 at 1/100 staining rat uterine tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)

AAA30956 at 1/100 staining human liver tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)

AAA30956 at 1/100 staining human heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)

AAA30956 at 1/100 staining mouse lung tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-MITF antibody

Description: MITF is a transcription factor that contains both basic helix-loop-helix and leucine zipper structural features. Plays a critical role in the differentiation of various cell types including neural crest-derived melanocytes, mast cells, osteoclasts and optic cup-derived retinal pigment epithelium.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557755
NCBI GeneID	4286
NCBI Accession #	NP_000239.1
NCBI GenBank Nucleotide #	NM_000248.3
UniProt Accession #	O75030; Q14841; Q9P2V0; Q9P2V1; Q9P2V2; Q9P2Y8; B4DJL2; D3K197; E9PFN0
Molecular Weight	52 kDa
NCBI Official Full Name	microphthalmia-associated transcription factor isoform 4
NCBI Official Synonym Full Names	melanocyte inducing transcription factor
NCBI Official Symbol	MITF
NCBI Official Synonym Symbols	MI; WS2; CMM8; WS2A; COMMAD; bHLHe32
NCBI Protein Information	microphthalmia-associated transcription factor
UniProt Protein Name	Microphthalmia-associated transcription factor
UniProt Gene Name	MITF
UniProt Synonym Gene Names	BHLHE32; bHLHe32

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mitf-antibody-western-blot-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/30956>