

Rabbit c-Jun Polyclonal Antibody | anti-JUN antibody

Phospho-c-Jun (Thr239) Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30982	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunoprecipitation, ELISA

PRODUCT SYNONYMS

c-Jun; Polyclonal Antibody; Phospho-c-Jun (Thr239) Antibody; Activator protein 1; AP 1; AP1; cJun; Enhancer Binding Protein AP1; Jun Activation Domain Binding Protein; JUN; Jun oncogene; JUN protein; Jun proto oncogene; JUN_HUMAN; JUNC; Oncogene JUN; p39; Proto oncogene c jun; Proto oncogene cJun; Proto-oncogene c-jun; Transcription Factor AP 1; Transcription factor AP-1; Transcription Factor AP1; V jun avian sarcoma virus 17 oncogene homolog; V jun sarcoma virus 17 oncogene homolog (avian); V jun sarcoma virus 17 oncogene homolog; V-jun avian sarcoma virus 17 oncogene homolog; vJun Avian Sarcoma Virus 17 Oncogene Homolog; anti-JUN antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-c-Jun (Thr239) antibody detects endogenous levels of c-Jun only when phosphorylated at Threonine 239
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	331
Applicable Applications for anti-JUN antibody	WB (Western Blot), IHC (Immunohistochemistry), IP (Immunoprecipitation), ELISA
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200
Immunogen	A synthesized peptide derived from human c-Jun around the phosphorylation site of Threonine 239
Subcellular Location	Nucleus.
Conjugation	Unconjugated
Epitope	Phospho Thr239
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




WB (Western Blot)

Western blot analysis of c-Jun phosphorylation expression in UV treated HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.




WB (Western Blot)

Western blot analysis of Phospho-c-Jun (Thr239) expression in various lysates




IHC (Immunohistochemistry)

AAA30982 at 1/100 staining rat intestinal tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)

AAA30982 at 1/100 staining rat gastric tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)

AAA30982 at 1/100 staining rat brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)

AAA30982 at 1/100 staining human TB tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-JUN antibody	Description: This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. Function: Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306). Subunit Structure: Heterodimer with either FOS or BATF3 or ATF7 (PubMed:10376527, PubMed:12087103, PubMed:15467742). The ATF7/JUN heterodimer is essential for ATF7 transactivation activity (PubMed:10376527). Interacts with DSIPI; the interaction inhibits the binding of active AP1 to its target DNA (By similarity). Interacts with HIVEP3 and MYBBP1A (By similarity). Interacts with SP1, SPIB and TCF20 (PubMed:10196196, PubMed:16478997, PubMed:8663478). Interacts with COPS5; the interaction leads indirectly to its phosphorylation (PubMed:8837781). Component of the SMAD3/SMAD4/JUN/FOS/complex which forms at the AP1 promoter site (PubMed:10995748). The SMAD3/SMAD4 heterodimer acts synergistically with the JUN/FOS heterodimer to activate transcription in response to TGF-beta (PubMed:9732876). Interacts (via its basic DNA binding and leucine zipper domains) with SMAD3 (via an N-terminal domain); the interaction is required for TGF-beta-mediated transactivation of the SMAD3/SMAD4/JUN/FOS/complex (PubMed:10995748). Interacts with methylated RNF187 (PubMed:20852630, PubMed:23624934). Binds to HIPK3. Interacts (when phosphorylated) with FBXW7 (PubMed:14739463). Found in a complex with PRR7 and FBXW7 (PubMed:27458189). Interacts with PRR7 and FBXW7; the interaction inhibits ubiquitination-mediated JUN degradation promoting its phosphorylation and transcriptional activity (PubMed:27458189). Post-translational Modifications: Ubiquitinated by the SCF(FBXW7), leading to its degradation (PubMed:14739463, PubMed:27458189). Ubiquitination takes place following phosphorylation, that promotes interaction with FBXW7 (PubMed:14739463). Phosphorylated by CaMK4 and PRKDC; phosphorylation enhances the transcriptional activity. Phosphorylated by HIPK3. Phosphorylated by DYRK2 at Ser-243; this primes the protein for subsequent phosphorylation by GSK3B at Thr-239. Phosphorylated at Thr-239, Ser-243 and Ser-249 by GSK3B; phosphorylation reduces its ability to bind DNA. Phosphorylated by PAK2 at Thr-2, Thr-8, Thr-89, Thr-93 and Thr-286 thereby promoting JUN-mediated cell proliferation and transformation. Phosphorylated by PLK3 following hypoxia or UV irradiation, leading to increase DNA-binding activity. Acetylated at Lys-271 by EP300. Similarity: Belongs to the bZIP family. Jun subfamily.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758616
NCBI GeneID	3725
NCBI Accession #	NP_002219.1
NCBI GenBank Nucleotide #	NM_002228.3
UniProt Accession #	P05412; Q6FHM7; Q96G93
Molecular Weight	Observed: 37 kDa Predicted: 36 kDa
NCBI Official Full Name	transcription factor AP-1
NCBI Official Synonym Full Names	Jun proto-oncogene, AP-1 transcription factor subunit
NCBI Official Symbol	JUN
NCBI Official Synonym Symbols	AP1; p39; AP-1; c-Jun
NCBI Protein Information	transcription factor AP-1
UniProt Protein Name	Transcription factor AP-1
UniProt Gene Name	JUN
UniProt Synonym Gene Names	AP1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-jun-antibody-western-blot-immunohistochemistry-immunoprecipitation-elisa/30982>