

Rabbit Cofilin Polyclonal Antibody | anti-CFL1 antibody

Phospho-Cofilin (Ser3) Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31025	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA

PRODUCT SYNONYMS


Cofilin; Polyclonal Antibody; Phospho-Cofilin (Ser3) Antibody; 18 kDa phosphoprotein; CFL 1; CFL; CFL1; COF1_HUMAN; Cofilin 1; Cofilin 1 non muscle; Cofilin non muscle isoform; Cofilin-1; epididymis secretory protein Li 15; HEL-S-15; non-muscle isoform; p18; anti-CFL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-Cofilin (Ser3) antibody detects endogenous levels of Cofilin only when phosphorylated at Serine 3
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	166
Applicable Applications for anti-CFL1 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB: 1:500-1:2000 IHC: 1/200 IF/ICC: 1:200
Immunogen	A synthesized peptide derived from human Cofilin around the phosphorylation site of Serine 3
Subcellular Location	Nucleus Matrix. Cytoplasm > Cytoskeleton. Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide.
Tissue Specificity	Widely distributed in various tissues.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Dog
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Dog (100%)
Conjugation	Unconjugated
Epitope	Phospho Ser3
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat ovarian tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat gastric tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat uterine tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31025 at 1/100 staining rat heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-CFL1 antibody	Description: Cofilin 1 a cytoskeletal protein that controls actin depolymerization. Has a 5-10-fold higher affinity for ATP-actin monomers than cofilin-1. May promote filament assembly rather than disassembly. Two alternatively spliced variants are described. Function: Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity. Regulates actin cytoskeleton dynamics. Important for normal progress through mitosis and normal cytokinesis. Plays a role in the regulation of cell morphology and cytoskeletal organization. Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:11812157, PubMed:15580268, PubMed:21834987, PubMed:23633677). Required for neural tube morphogenesis and neural crest cell migration (By similarity). Subunit Structure: Can bind G-and F-actin in a 1:1 ratio of cofilin to actin. It is a major component of intranuclear and cytoplasmic actin rods. Post-translational Modifications: Inactivated by phosphorylation on Ser-3. Phosphorylated on Ser-3 in resting cells (By similarity). Dephosphorylated by PDXP/chronophin; this restores its activity in promoting actin filament depolymerization. The phosphorylation of Ser-24 may prevent recognition of the nuclear localization signal (By similarity). Phosphorylated via a ARRB1-RAC1-LIMK1-PAK1 cascade upon active ligand stimulation of atypical chemokine receptor ACKR2. Similarity: Belongs to the actin-binding proteins ADF family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	5031635
NCBI GeneID	1072
NCBI Accession #	NP_005498.1
NCBI GenBank Nucleotide #	NM_005507.2
UniProt Accession #	P23528; Q53Y87; Q9UCA2; B3KUQ1
Molecular Weight	Observed: 19 kDa Predicted: 19 kDa
NCBI Official Full Name	cofilin-1
NCBI Official Synonym Full Names	cofilin 1
NCBI Official Symbol	CFL1
NCBI Official Synonym Symbols	CFL; cofilin; HEL-S-15
NCBI Protein Information	cofilin-1
UniProt Protein Name	Cofilin-1
UniProt Gene Name	CFL1
UniProt Synonym Gene Names	CFL; p18

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.