

Rabbit anti-Human IGHA1 Polyclonal Antibody | anti-IGHA1 antibody

IGHA1 Antibody

CATALOG # AAA310529	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

IGHA1; Polyclonal Antibody; IGHA1 Antibody; IgA1; anti-IGHA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	The antibody detects endogenous levels of total IGHA1 protein.
Purity/Purification	Antigen affinity purification
Form/Format	pH 7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Concentration	0.9 mg/mL (varies by lot)
Sequence Length	353
Applicable Applications for anti-IGHA1 antibody	IHC (Immunohistochemistry)
Immunogen	Fusion protein of human IGHA1
Immunogen Type	Peptide
Preparation and Storage	Store at -20 degree C.

IMAGES



IHC (Immunohistochemistry)

The image is immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using 47254(IGHA1 Antibody) at dilution 1/50.(Original magnification: 200



IHC (Immunohistochemistry)

The image is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using 47254(IGHA1 Antibody) at dilution 1/50.(Original magnification: 200

ADDITIONAL INFORMATION

Related Product Information for anti-IGHA1 antibody	Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Ig alpha is the major immunoglobulin class in body secretions (PubMed:2241915).
Product Categories/Family for anti-IGHA1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	113584
NCBI GeneID	3493
UniProt Accession #	P01876
Molecular Weight	37,655 Da
NCBI Official Full Name	Immunoglobulin heavy constant alpha 1
NCBI Official Synonym Full Names	immunoglobulin heavy constant alpha 1
NCBI Official Symbol	IGHA1
NCBI Official Synonym Symbols	IgA1
UniProt Protein Name	Immunoglobulin heavy constant alpha 1
UniProt Gene Name	IGHA1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/igha1-antibody-immunohistochemistry-elisa/310529>