

Rabbit anti-Human SIRT1 Monoclonal Antibody | anti-SIRT1 antibody

SIRT1 (Phospho-T530) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA311039	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

SIRT1; Monoclonal Antibody; SIRT1 (Phospho-T530) Antibody; 75SirT1 antibody; hSIR2 antibody; hSIRT1 antibody; HST2; S. cerevisiae; homolog of antibody; NAD dependent deacetylase sirtuin 1 antibody; NAD dependent protein deacetylase sirtuin 1 antibody; OTTHUMP00000198111 antibody; OTTHUMP00000198112 antibody; Regulatory protein SIR2 homolog 1 antibody; SIR1_HUMAN antibody; SIR2 antibody; SIR2 like 1 antibody; SIR2 like protein 1 antibody; SIR2; S.cerevisiae; homolog-like 1 antibody; SIR2-like protein 1 antibody; SIR2ALPHA antibody; SIR2L1 antibody; Sirt1 antibody; SirtT1 75 kDa fragment antibody; Sirtuin (silent mating type information regulation 2 homolog) 1 (S. cerevisiae) antibody; Sirtuin 1 antibody; Sirtuin type 1 antibody; anti-SIRT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Clone Number	JJ206-6
Purity/Purification	ProA affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-SIRT1 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Antibody Type	Recombinant Antibody
Immunogen	Synthetic phospho-peptide corresponding to residues surrounding Thr530 of human SIRT1.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of 293 cells with Phospho-SIRT1 (T530) antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




ICC (Immunocytochemistry)

ICC staining Phospho-SIRT1 (T530) in 293 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




ICC (Immunocytochemistry)

ICC staining Phospho-SIRT1 (T530) in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC (Immunocytochemistry)

ICC staining Phospho-SIRT1 (T530) in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-Phospho-SIRT1 (T530) antibody. Counter stained with hematoxylin.

ADDITIONAL INFORMATION

Related Product Information for anti-SIRT1 antibody	The silent information regulator (SIR2) family of genes are highly-conserved from prokaryotes to eukaryotes and are involved in diverse processes, including transcriptional regulation, cell cycle progression, DNA-damage repair and aging. In <i>S. cerevisiae</i> , Sir2p deacetylates histones in an NAD-dependent manner, which regulates silencing at the telomeric, rDNA and silent mating-type loci. Sir2p is the founding member of a large family, designated sirtuins, which contain a conserved catalytic domain. The human homologs, which include SIRT1-7, are divided into four main branches: SIRT1-3 are class I, SIRT4 is class II, SIRT5 is class III and SIRT6-7 are class IV. SIRT1 has the closest homology to the yeast Sir2p and is widely expressed in fetal and adult tissues. SIRT1 is highly expressed in heart, brain and skeletal muscle, with low expression in lung and placenta. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lys 382.
Product Categories/Family for anti-SIRT1 antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	215982798
NCBI GeneID	23411
NCBI Accession #	NP_001135970.1
NCBI GenBank Nucleotide #	NM_001142498.1
UniProt Accession #	Q96EB6 ; Q2XNF6 ; Q5JYQ0 ; Q9GZR9 ; Q9Y6F0
Molecular Weight	Calculated MW: 82kDa
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-1 isoform b
NCBI Official Synonym Full Names	sirtuin 1
NCBI Official Symbol	SIRT1
NCBI Official Synonym Symbols	SIR2L1

NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-1; hSIR2; hSIRT1; SIR2alpha; sir2-like 1; sirtuin type 1; SIR2-like protein 1; regulatory protein SIR2 homolog 1; NAD-dependent deacetylase sirtuin-1
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-1
UniProt Gene Name	SIRT1
UniProt Synonym Gene Names	SIR2L1; hSIRT1; hSIR2; 75SirT1
UniProt Entry Name	SIR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/sirt1-antibody-clone-jj206-6-flow-cytometry-immunohistochemistry-immunofluorescence-immunocytochemistry-western-blot-elisa/311039>