

Rabbit CALR Polyclonal Antibody | anti-CALR antibody

CALR Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31108	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA

PRODUCT SYNONYMS


CALR; Polyclonal Antibody; CALR Antibody; Autoantigen RO; CALR protein; CALR_HUMAN; Calregulin; Calreticulin; cC1qR; CRP55; CRT; CRTC; Endoplasmic reticulum resident protein 60; Epididymis secretory sperm binding protein Li 99n; ERp60; FLJ26680; grp60; HACBP; HEL S 99n; RO; Sicca syndrome antigen A (autoantigen Ro; calreticulin); Sicca syndrome antigen A; SSA; anti-CALR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	CALR antibody detects endogenous levels of total CALR
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	417
Applicable Applications for anti-CALR antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB: 1:500-1:1000 IHC: 1:50-1:200 IF/ICC: 1:100-1:500
Immunogen	A synthesized peptide
Subcellular Location	Endoplasmic Reticulum Lumen. Cytoplasm > Cytosol. Secreted > Extracellular Space > Extracellular Matrix. Cell Surface. Also found in cell surface (T cells), cytosol and extracellular matrix. Associated with the lytic granules in the cytolytic T-lymphocytes.
Predicted Cross Reactivity	Pig, Zebrafish, Bovine, Horse, Sheep, Rabbit, Xenopus
Similarity	Pig (91%), Zebrafish (91%), Bovine (91%), Horse (91%), Sheep (91%), Rabbit (91%), Xenopus (91%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




WB (Western Blot)
Western blot analysis of extracts from HepG2 and 293, using CALR Antibody.




WB (Western Blot)
Western blot analysis of extracts from COS-7 cells, using CALR antibody. The lane on the left is treated with the antigen-specific peptide.




WB (Western Blot)
Western blot analysis of CALR expression in Mouse muscle lysate



IHC (Immunohistochemistry)
AAA31108 at 1/100 staining Human prostate tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IF (Immunofluorescence)
AAA31108 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from mouse brain, using CALR Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CALR antibody	Function: Calcium-binding chaperone that promotes folding, oligomeric assembly and quality control in the endoplasmic reticulum (ER) via the calreticulin/calnexin cycle. This lectin interacts transiently with almost all of the monoglucosylated glycoproteins that are synthesized in the ER. Interacts with the DNA-binding domain of NR3C1 and mediates its nuclear export. Involved in maternal gene expression regulation. May participate in oocyte maturation via the regulation of calcium homeostasis (By similarity). Subunit Structure: Monomer. Component of an EIF2 complex at least composed of CELF1/CUGBP1, CALR, CALR3, EIF2S1, EIF2S2, HSP90B1 and HSPA5. Interacts with PDIA3/Erp57 and SPACA9 (By similarity). Interacts with TRIM21 (PubMed:8666824). Interacts with NR3C1 (PubMed:11149926). Interacts with PPIB (PubMed:20801878). Interacts (via P-domain) with PDIA5 (PubMed:23614004). Interacts with GABARAP (PubMed:19154346). Similarity: Can be divided into a N-terminal globular domain, a proline-rich P-domain forming an elongated arm-like structure and a C-terminal acidic domain. The P-domain binds one molecule of calcium with high affinity, whereas the acidic C-domain binds multiple calcium ions with low affinity. The interaction with glycans occurs through a binding site in the globular lectin domain. The zinc binding sites are localized to the N-domain. Associates with PDIA3 through the tip of the extended arm formed by the P-domain. Belongs to the calreticulin family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757900
NCBI GeneID	811
NCBI Accession #	NP_004334.1
NCBI GenBank Nucleotide #	NM_004343.3
UniProt Accession #	P27797; Q6IAT4; Q9UDG2
Molecular Weight	Observed: 48 kD Predicted: 49 kDa
NCBI Official Full Name	calreticulin
NCBI Official Synonym Full Names	calreticulin
NCBI Official Symbol	CALR
NCBI Official Synonym Symbols	RO; CRT; SSA; cC1qR; HEL-S-99n
NCBI Protein Information	calreticulin
UniProt Protein Name	Calreticulin
UniProt Gene Name	CALR
UniProt Synonym Gene Names	CRTC; ERp60

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/calr-antibody-western-blot-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/31108>