

Rabbit EFNB2 Polyclonal Antibody | anti-EFNB2 antibody

EFNB2 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31132	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA

PRODUCT SYNONYMS


EFNB2; Polyclonal Antibody; EFNB2 Antibody; EFN B2; EFNB 2; Efnb2; EFNB2_HUMAN; Eph related receptor tyrosine kinase ligand 5; EPH-related receptor tyrosine kinase ligand 5; ephrin B2; Ephrin-B2; EphrinB2; EPLG 5; EPLG5; Htk L; HTK ligand; HTK-L; HTKL; LERK 5; LERK-5; LERK5; Ligand of eph related kinase 5; MGC126226; MGC126227; MGC126228; OTTMUSP00000024973; anti-EFNB2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	EFNB2 antibody detects endogenous levels of total EFNB2
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	333
Applicable Applications for anti-EFNB2 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 ICC/IF: 1:100-500
Immunogen	A synthesized peptide derived from human EFNB2
Subcellular Location	Membrane.
Tissue Specificity	Lung and kidney.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IHC (Immunohistochemistry)
AAA31132 at 1/100 staining rat testicular tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)
AAA31132 at 1/100 staining rat liver tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.




IHC (Immunohistochemistry)
AAA31132 at 1/100 staining rat lung tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31132 at 1/100 staining rat Intestinal tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31132 at 1/200 staining human myosarcoma tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)
AAA31132 at 1/200 staining human seminoma tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-EFNB2 antibody	Description: This gene encodes a member of the ephrin (EPH) family. The ephrins and EPH-related receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been implicated in mediating developmental events, especially in the nervous system and in erythropoiesis. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. This gene encodes an EFNB class ephrin which binds to the EPHB4 and EPHA3 receptors. Function: Cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Binds to receptor tyrosine kinase including EPHA4, EPHA3 and EPHB4. Together with EPHB4 plays a central role in heart morphogenesis and angiogenesis through regulation of cell adhesion and cell migration. EPHB4-mediated forward signaling controls cellular repulsion and segregation from EFNB2-expressing cells. May play a role in constraining the orientation of longitudinally projecting axons. Subunit Structure: Interacts with PDZRN3 (By similarity). Binds to the receptor tyrosine kinases EPHA4, EPHB4 and EPHA3. Post-translational Modifications: Inducible phosphorylation of tyrosine residues in the cytoplasmic domain. Similarity: Belongs to the ephrin family.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758250
NCBI GeneID	1948
NCBI Accession #	NP_004084.1
NCBI GenBank Nucleotide #	NM_004093.3
UniProt Accession #	P52799; Q5JV56
Molecular Weight	Observed: 36 kDa Predicted: 37 kDa
NCBI Official Full Name	ephrin-B2
UniProt Protein Name	Ephrin-B2
UniProt Gene Name	EFNB2
UniProt Synonym Gene Names	EPLG5; HTKL; LERK5; LERK-5; HTK-L

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/efnb2-antibody-western-blot-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/31132>