

Rabbit P21 Polyclonal Antibody | anti-P21 antibody

P21 Polyclonal Antibody

CATALOG # AAA311324	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

P21; Polyclonal Antibody; P21 Polyclonal Antibody; CDKN1A; CAP20; CDKN1; CIP1; MDA-6; SDI1; WAF1; p21CIP1; cyclin-dependent kinase inhibitor 1; anti-P21 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 50% glycerol, pH7.4
Applicable Applications for anti-P21 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human P21 (NP_001207707.1).
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using P21 antibody.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human uterine cancer using P21 antibody.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon carcinoma using P21 antibody.



WB (Western Blot)

Western blot analysis of extracts from normal (control) and P21 knockout (KO) HeLa cells, using P21 antibody.



WB (Western Blot)

Western blot analysis of extracts of U-87MG cells, using P21 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-P21 antibody

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.

Product Categories/Family for anti-P21 antibody

Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	334085242
NCBI GeneID	1026
NCBI Accession #	NP_001207707.1
NCBI GenBank Nucleotide #	NM_001220778.1
UniProt Accession #	P38936; Q14010; Q9BUT4
Molecular Weight	~47kD

NCBI Official Full Name	cyclin-dependent kinase inhibitor 1
NCBI Official Synonym Full Names	cyclin-dependent kinase inhibitor 1A (p21, Cip1)
NCBI Official Symbol	CDKN1A
NCBI Official Synonym Symbols	P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1
NCBI Protein Information	cyclin-dependent kinase inhibitor 1; DNA synthesis inhibitor; CDK-interacting protein 1; CDK-interaction protein 1; wild-type p53-activated fragment 1; melanoma differentiation associated protein 6
UniProt Protein Name	Cyclin-dependent kinase inhibitor 1
UniProt Gene Name	CDKN1A
UniProt Synonym Gene Names	CAP20; CDKN1; CIP1; MDA6; PIC1; SDI1; WAF1; MDA-6
UniProt Entry Name	CDN1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/p21-antibody-immunohistochemistry-western-blot-elisa/311324>