

Rabbit SHP-2 Polyclonal Antibody | anti-SHP-2 antibody

Phospho-SHP-2 (Tyr62) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31155	Rabbit	Human, Mouse, Rat Predicted: Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus	Western Blot, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SHP-2; Polyclonal Antibody; Phospho-SHP-2 (Tyr62) Antibody; BTP3; CFC; JMML; METCDS; MGC14433; NS1; OTTHUMP00000166107; OTTHUMP00000166108; Protein tyrosine phosphatase 2; Protein tyrosine phosphatase 2C; Protein tyrosine phosphatase non receptor type 11; Protein-tyrosine phosphatase 1D; Protein-tyrosine phosphatase 2C; PTN11_HUMAN; PTP-1D; PTP-2C; PTP1D; PTP2C; PTPN11; SAP2; SH-PTP2; SH-PTP3; SH2 domain containing protein tyrosine phosphatase 2; SHP 2; Shp2; SHPTP2; SHPTP3; Syp; Tyrosine-protein phosphatase non-receptor type 11; anti-SHP-2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted: Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-SHP-2 (Tyr62) Antibody detects endogenous levels of SHP-2 only when phosphorylated at Tyr62.
Purity/Purification	Purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form/Format	Liquid; Rabbit IgG in Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	593
Applicable Applications for anti-SHP-2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA (Peptide ELISA)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 ELISA(peptide): 1:20000-1:40000 IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.
Immunogen	A synthesized peptide derived from human SHP-2 around the phosphorylation site of Tyr62.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western blot analysis of extracts from A431, using Phospho-SHP-2 (Tyr62) Antibody. Lane1 was treated with phospho-blocking peptide, Lane2 was treated with non-phospho-blocking peptide.



WB (Western Blot)

Western blot analysis of Phospho-SHP-2 (Tyr62) in lysates of A431, using Phospho-SHP-2 (Tyr62) Antibody.




IHC (Immunohistochemistry)

At 1/100 staining human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary




IHC (Immunohistochemistry)

At 1/100 staining rat brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary



IHC (Immunohistochemistry)

At 1/100 staining rat heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary



IHC (Immunohistochemistry)

At 1/100 staining mouse heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary

ADDITIONAL INFORMATION

Related Product Information for anti-SHP-2 antibody

Description: Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus. Positively regulates MAPK signal transduction pathway (PubMed:28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity. Dephosphorylates CDC73 (PubMed:26742426).

Post Translational Modifications: Phosphorylated on Tyr-546 and Tyr-584 upon receptor protein tyrosine kinase activation; which creates a binding site for GRB2 and other SH2-containing proteins. Phosphorylated upon activation of the receptor-type kinase FLT3. Phosphorylated upon activation of the receptor-type kinase PDGFRA (By similarity). Phosphorylated by activated PDGFRB.

Subunit Structure: Interacts with phosphorylated LIME1 and BCAR3. Interacts with SHB and INPP5D/SHIP1 (By similarity). Interacts with MILR1 (tyrosine-phosphorylated). Interacts with FLT1 (tyrosine-phosphorylated), FLT3 (tyrosine-phosphorylated), FLT4 (tyrosine-phosphorylated), KIT and GRB2. Interacts with PDGFRA (tyrosine phosphorylated). Interacts (via SH2 domain) with TEK/TIE2 (tyrosine phosphorylated) (By similarity). Interacts with PTPNS1 and CD84. Interacts with phosphorylated SIT1 and MPZL1. Interacts with FCRL4, FCRL6 and ANKHD1. Interacts with KIR2DL1; the interaction is enhanced by ARRB2. Interacts with GAB2. Interacts with TERT; the interaction retains TERT in the nucleus. Interacts with PECAM1 and FER. Interacts with EPHA2 (activated); participates in PTK2/FAK1 dephosphorylation in EPHA2 downstream signaling. Interacts with ROS1; mediates PTPN11 phosphorylation. Interacts with PDGFRB (tyrosine phosphorylated); this interaction increases the PTPN11 phosphatase activity. Interacts with GAREM1 isoform 1 (tyrosine phosphorylated); the interaction increases MAPK/ERK activity and does not affect the GRB2/SOS complex formation. Interacts with CDC73 (PubMed:26742426). Interacts with CEACAM1 (via cytoplasmic domain); this interaction depends on the monomer/dimer equilibrium and is phosphorylation-dependent (By similarity). Interacts with MPIG6B (via ITIM motif) (PubMed:23112346). Interacts with SIGLEC10 (By similarity). Interacts with FCRL3 (via phosphorylated ITIM motifs) (PubMed:11162587, PubMed:19843936).

Similarity: The SH2 domains repress phosphatase activity. Binding of these domains to phosphotyrosine-containing proteins relieves this auto-inhibition, possibly by inducing a conformational change in the enzyme. Belongs to the protein-tyrosine phosphatase family. Non-receptor class 2 subfamily.

NCBI AND UNIPROT INFORMATION

NCBI GI #	33356177
NCBI GeneID	5781
NCBI Accession #	NP_002825.3
NCBI GenBank Nucleotide #	NP_002825.3

UniProt Accession #	Q06124
Molecular Weight	68.4 kDa
NCBI Official Full Name	tyrosine-protein phosphatase non-receptor type 11 isoform 1
NCBI Official Synonym Full Names	protein tyrosine phosphatase non-receptor type 11
NCBI Official Symbol	PTPN11
NCBI Official Synonym Symbols	CFC; NS1; JMML; SHP2; BPTP3; PTP2C; METCDS; PTP-1D; SH-PTP2; SH-PTP3
NCBI Protein Information	tyrosine-protein phosphatase non-receptor type 11
UniProt Protein Name	Tyrosine-protein phosphatase non-receptor type 11
UniProt Gene Name	PTPN11
UniProt Synonym Gene Names	PTP2C; SHPTP2; PTP-1D; PTP-2C; SHP-2; Shp2
UniProt Entry Name	PTN11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/shp-2-antibody-western-blot-immunohistochemistry-peptide-elisa-elisa/31155>