

Rabbit Clathrin Polyclonal Antibody | anti-CLTC antibody

Clathrin Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31165	Rabbit	Human, Mouse, Rat	Western Blot, Immunofluorescence, Immunocytochemistry, ELISA

PRODUCT SYNONYMS

Clathrin; Polyclonal Antibody; Clathrin Antibody; Adapter related protein complex 1 beta 1 subunit; Adaptor Protein Complex AP1 Beta 1 Subunit; ADTB1; AP 1 complex subunit beta 1; AP105A; AP1B1; BAM22; Beta Adaptin 1; Beta Prime Adaptin; CLAPB2; Clathrin Assembly Protein Complex 1 Beta Large Chain; Clathrin heavy chain 1; CLH 17; CLH17; CLTC; CLTCL2; Golgi Adaptor HA1/AP1 Adaptin Beta Subunit; KIAA0034; Plasma Membrane Adaptor HA2/AP2 Adaptor Beta Subunit; anti-CLTC antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Clathrin Antibody detects endogenous levels of total Clathrin.
Purity/Purification	Peptide affinity purification
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Sequence Length	1675
Applicable Applications for anti-CLTC antibody	WB (Western Blot), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB: 1:500-1:2000 IF: 1:100-1:500 ICC: 1:100-1:500 IHC: 1:50-1:200 ELISA(peptide): 1:20,000-1:40,000
Immunogen	A synthesized peptide derived from human Clathrin, corresponding to a region within C-terminal amino acids.
Fragment	Fab fragment
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IF (Immunofluorescence)

Staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF13215 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31165 at 1/100 staining Human pancreatic cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31165 at 1/100 staining Human kidney cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31165 at 1/100 staining Human kidney cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31165 at 1/100 staining Human ovarian cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31165 at 1/100 staining Mouse spleen tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CLTC antibody	Description: Clathrin is the major protein of the polyhedral coat of coated pits and vesicles. Two different adapter protein complexes link the clathrin lattice either to the plasma membrane or to the trans-Golgi network. Acts as component of the TACC3/ch-TOG/clathrin complex proposed to contribute to stabilization of kinetochore fibers of the mitotic spindle by acting as inter-microtubule bridge (PubMed:15858577, PubMed:16968737, PubMed:21297582). The TACC3/ch-TOG/clathrin complex is required for the maintenance of kinetochore fiber tension (PubMed:23532825). Plays a role in early autophagosome formation (PubMed:20639872).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758012
NCBI GeneID	1213
NCBI Accession #	NP_004850.1
NCBI GenBank Nucleotide #	NP_004850.1
UniProt Accession #	Q00610
Molecular Weight	191kDa
NCBI Official Full Name	clathrin heavy chain 1 isoform 1
NCBI Official Synonym Full Names	clathrin heavy chain
NCBI Official Symbol	CLTC
NCBI Official Synonym Symbols	Hc; CHC; CHC17; MRD56; CLH-17; CLTCL2
NCBI Protein Information	clathrin heavy chain 1
UniProt Protein Name	Clathrin heavy chain 1
UniProt Gene Name	CLTC
UniProt Synonym Gene Names	CLH17; CLTCL2; KIAA0034; CLH-17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/clathrin-antibody-western-blot-immunofluorescence-immunocytochemistry-elisa/31165>