

Rabbit anti-Human, Mouse Cadherin EGF LAG Seven-Pass G-Type 2 Polyclonal Antibody | anti-CELSR2 antibody

Cadherin EGF LAG Seven-Pass G-Type Receptor 2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA311874	Rabbit	Human, Mouse	Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

Cadherin EGF LAG Seven-Pass G-Type 2; Polyclonal Antibody; Cadherin EGF LAG Seven-Pass G-Type Receptor 2 Antibody; Cadherin EGF LAG seven pass G type receptor 2 flamingo Drosophila homolog antibody; Cadherin EGF LAG seven-pass G-type receptor 2 antibody; Cadherin family member 10 antibody; CDHF10 antibody; CELR2_HUMAN antibody; Celsr2 antibody; EGF like domain multiple 2 antibody; EGF-like protein 2 antibody; EGFL2 antibody; epidermal growth factor like 2 antibody; Epidermal growth factor-like protein 2 antibody; Flamingo homolog 3 antibody; Flamingo1 antibody; KIAA0279 antibody; MEGF3 antibody; Multiple EGF-like domains protein 3 antibody; Multiple epidermal growth factor-like domains 3 antibody; Multiple epidermal growth factor-like domains protein 3 antibody; anti-CELSR2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-CELSR2 antibody	ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	Recombinant protein within human Cadherin EGF LAG seven-pass G-type receptor 2 1-250 aa.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.

IMAGES



ICC (Immunocytochemistry)

ICC image of CELSR2 antibody stained D3 (Murine Embryonic Stem Cell Line).The secondary antibody (red) was goat anti-rabbit IgG (H+L) alexa555 conjugated.



WB (Western Blot)

Western blot analysis on recombinant protein of CELSR2 using anti- CELSR2 polyclonal antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CELSR2 antibody	Cadherin EGF LAG seven-pass G-type receptor 2 belongs to the G-protein coupled receptor 2 family, highest expressed in brain and testis. The protein encoded by the gene CELSR2 is a member of the flamingo subfamily, part of the cadherin superfamily. It may have an important role in cell/cell signaling during nervous system formation.
Product Categories/Family for anti-CELSR2 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	13325064
NCBI GeneID	1952
NCBI Accession #	NP_001399.1
NCBI GenBank Nucleotide #	NM_001408.2
UniProt Accession #	Q9HCU4; Q5T2Y7; Q92566
Molecular Weight	317,453 Da
NCBI Official Full Name	cadherin EGF LAG seven-pass G-type receptor 2
NCBI Official Synonym Full Names	cadherin, EGF LAG seven-pass G-type receptor 2
NCBI Official Symbol	CELSR2
NCBI Official Synonym Symbols	EGFL2; MEGF3; CDHF10; Flamingo1
NCBI Protein Information	cadherin EGF LAG seven-pass G-type receptor 2; EGF-like protein 2; flamingo homolog 3; cadherin family member 10; EGF-like-domain, multiple 2; epidermal growth factor-like 2; multiple EGF-like domains protein 3; epidermal growth factor-like protein 2; mul
UniProt Protein Name	Cadherin EGF LAG seven-pass G-type receptor 2
UniProt Gene Name	CELSR2
UniProt Synonym Gene Names	CDHF10; EGFL2; KIAA0279; MEGF3; EGF-like protein 2
UniProt Entry Name	CELR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cadherin-egf-lag-seven-pass-g-type-2-antibody-immunocytochemistry-western-blot-elisa/311874>