

Rabbit KGF Polyclonal Antibody | anti-FGF7 antibody

KGF Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31200	Rabbit	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Zebrafish(100%), Bovine(100%), Horse(100%), Sheep(100%), Rabbit(100%), Dog(100%), Chicken(100%), Xenopus(100%)	Immunofluorescence, Immunocytochemistry, ELISA

PRODUCT SYNONYMS

KGF; Polyclonal Antibody; KGF Antibody; FGF 7; FGF-7; Fgf7; FGF7_HUMAN; Fibroblast growth factor 7; HBGF 7; HBGF-7; HBGF7; Heparin binding growth factor 7; Heparin-binding growth factor 7; Keratinocyte growth factor; anti-FGF7 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Zebrafish(100%), Bovine(100%), Horse(100%), Sheep(100%), Rabbit(100%), Dog(100%), Chicken(100%), Xenopus(100%)
Clonality	Polyclonal
Isotype	IgG
Specificity	KGF Antibody detects endogenous levels of total KGF.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Sequence Length	194
Applicable Applications for anti-FGF7 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	IF: 1:100-1:500 ICC: 1:100-1:500 WB: 1:500-1:2000 IHC: 1:50-1:200 ELISA(peptide): 1:20,000-1:40,000
Immunogen	A synthesized peptide derived from human KGF, corresponding to a region within the internal amino acids.
Fragment	Fab fragment
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IF (Immunofluorescence)

AAA31200 staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AAA31200 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody. The nuclear counter stain is DAPI(blue).




IHC (Immunohistochemistry-Paraffin)

AAA31200 at 1/100 staining Human esophageal cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31200 at 1/100 staining Human colorectal cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31200 at 1/100 staining Mouse kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31200 at 1/100 staining Rat kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31200 at 1/100 staining Rat colorectal tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-FGF7 antibody	Function: Plays an important role in the regulation of embryonic development, cell proliferation and cell differentiation. Required for normal branching morphogenesis. Growth factor active on keratinocytes. Possible major paracrine effector of normal epithelial cell proliferation. Subcellular Location: Secreted. Tissue Specificity: Epithelial cell. Subunit Structure: Interacts with FGFBP1. Interacts with FGFR2. Affinity between fibroblast growth factors (FGFs) and their receptors is increased by heparan sulfate glycosaminoglycans that function as coreceptors. Similarity: Belongs to the heparin-binding growth factors family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503705
NCBI GeneID	2252
NCBI Accession #	NP_002000.1
NCBI GenBank Nucleotide #	NP_002000.1
UniProt Accession #	P21781
Molecular Weight	Observed Molecular Weight: (Observed)22kD. Predicted Molecular Weight: (Calculated)23kDa.
NCBI Official Full Name	fibroblast growth factor 7
NCBI Official Synonym Full Names	fibroblast growth factor 7
NCBI Official Symbol	FGF7
NCBI Official Synonym Symbols	KGF; HBGF-7
NCBI Protein Information	fibroblast growth factor 7
UniProt Protein Name	Fibroblast growth factor 7
UniProt Gene Name	FGF7
UniProt Synonym Gene Names	KGF; FGF-7; HBGF-7
UniProt Entry Name	FGF7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.