

Rabbit Integrin alpha V Monoclonal Antibody | anti-ITGAV antibody

Integrin alpha V Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA312246	Rabbit	Human, Mouse, Rat	Flow Cytometry, Functional Assay, Western Blot


PRODUCT SYNONYMS

Integrin alpha V; Monoclonal Antibody; Integrin alpha V Antibody; CD51 antibody; Integrin alpha-V light chain antibody; integrin; alpha V (vitronectin receptor; alpha polypeptide; antigen CD51) antibody; Integrin; alpha V antibody; beta 6 antibody; ITAV_HUMAN antibody; ITGAV antibody; ITGB6 antibody; Vitronectin receptor subunit alpha antibody; anti-ITGAV antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Clone Number	SC56-07
Purity/Purification	ProA affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-ITGAV antibody	FCM/FACS (Flow Cytometry), WB (Western Blot)
Antibody Type	Recombinant Antibody
Immunogen	Recombinant protein
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.

IMAGES



FCM/FACS (Flow Cytometry)

Flow cytometric analysis of MCF-7 cells with Integrin alpha V antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.



WB (Western Blot)

Western blot analysis of Integrin alpha V+beta 6 on different lysates using anti-Integrin alpha V+beta 6 antibody at 1/1,000 dilution. Positive control: Lane 1: A549 Lane 2: PC-12

ADDITIONAL INFORMATION

Related Product Information for anti-ITGAV antibody	Integrins are heterodimers composed of noncovalently associated transmembrane a and b subunits. The 16 a and 8 b subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and vitronectin. Certain integrins can also bind to soluble ligands, such as fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.
Product Categories/Family for anti-ITGAV antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	223468595
NCBI GeneID	3685
NCBI Accession #	NP_001138471.1
NCBI GenBank Nucleotide #	NM_001144999.2
UniProt Accession #	P06756; Q53SK4; Q59EB7; Q6LD15; A0AV67; B0LPF4; B7Z883; D3DPG8; E7EWZ6
Molecular Weight	116,038 Da
NCBI Official Full Name	integrin alpha-V isoform 2
NCBI Official Synonym Full Names	integrin, alpha V
NCBI Official Symbol	ITGAV
NCBI Official Synonym Symbols	CD51; MSK8; VNRA; VTNR
NCBI Protein Information	integrin alpha-V; integrin alphaVbeta3; vitronectin receptor subunit alpha; antigen identified by monoclonal antibody L230; integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51)
UniProt Protein Name	Integrin alpha-V
UniProt Gene Name	ITGAV
UniProt Synonym Gene Names	MSK8; VNRA
UniProt Entry Name	ITAV_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.