

Rabbit LIMK1 Polyclonal Antibody | anti-LIMK1 antibody

Anti-LIMK1 (Ab-508) antibody

CATALOG # AAA312930	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

LIMK1; Polyclonal Antibody; Anti-LIMK1 (Ab-508) antibody; LIMK; LIMK-1; anti-LIMK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Antigen affinity purification
Form/Format	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Concentration	0.8mg/ml (varies by lot)
Applicable Applications for anti-LIMK1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthetic peptide of human LIMK1
Preparation and Storage	Store at -20 degree C

IMAGES



IHC (Immunohistochemistry)

The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using LIMK1(Ab-508) Antibody at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: x200)



SDS-PAGE

Gel: 8%SDS-PAGE Lysate: 40 ug Lane: HUVEC cell lysate Primary antibody: LIMK1(Ab-508) Antibody at dilution 1/350 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 30 seconds

ADDITIONAL INFORMATION

Related Product Information for anti-LIMK1 antibody	There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is a serine/threonine kinase that regulates actin polymerization via phosphorylation and inactivation of the actin binding factor cofilin. This protein is ubiquitously expressed during development and plays a role in many cellular processes associated with cytoskeletal structure. This protein also stimulates axon growth and may play a role in brain development. LIMK1 hemizyosity is implicated in the impaired visuospatial constructive cognition of Williams syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
Product Categories/Family for anti-LIMK1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	324715046
NCBI GeneID	3984
NCBI Accession #	NP_001191355.1
NCBI GenBank Nucleotide #	NM_001204426.1
UniProt Accession #	P53667; O15283; Q15820; Q15821; Q75MU3; Q9Y5Q1; B7Z6I8; D3DXF4; D3DXF5
Molecular Weight	72,585 Da
NCBI Official Full Name	LIM domain kinase 1 isoform 2
NCBI Official Synonym Full Names	LIM domain kinase 1
NCBI Official Symbol	LIMK1
NCBI Official Synonym Symbols	LIMK; LIMK-1
NCBI Protein Information	LIM domain kinase 1; LIM motif-containing protein kinase
UniProt Protein Name	LIM domain kinase 1
UniProt Gene Name	LIMK1
UniProt Synonym Gene Names	LIMK; LIMK-1
UniProt Entry Name	LIMK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/limk1-antibody-immunohistochemistry-western-blot-elisa/312930>