

Rabbit RSK1 p90 Polyclonal Antibody | anti-RSK1p90 antibody

Phospho-RSK1 p90 (Thr348) Antibody


CATALOG # AAA31316	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, ELISA
------------------------------	-----------------------	--	--

PRODUCT SYNONYMS

RSK1 p90; Polyclonal Antibody; Phospho-RSK1 p90 (Thr348) Antibody; 90kDa ribosomal protein S6 kinase 1; dJ590P13.1 (ribosomal protein S6 kinase; 90kD; polypeptide 1; dJ590P13.1; EC 2.7.11.1; HU 1; HU1; KS6A1_HUMAN; MAP kinase activated protein kinase 1a; MAP kinase-activated protein kinase 1a; MAPK-activated protein kinase 1a; MAPKAP kinase 1a; MAPKAPK-1a; MAPKAPK1A; MGC79981; Mitogen-activated protein kinase-activated protein kinase 1A; OTTHUMP00000004113; p90 RSK1; p90-RSK 1; p90rsk; p90RSK1; p90S6K; pp90RSK1; Ribosomal protein S6 kinase 90kD 1; Ribosomal protein S6 kinase 90kD polypeptide 1; Ribosomal protein S6 kinase 90kDa polypeptide 1; Ribosomal protein S6 kinase alpha 1; Ribosomal protein S6 kinase alpha-1; Ribosomal protein S6 kinase polypeptide 1; Ribosomal S6 kinase 1; RPS6K1 alpha; rps6ka; Rps6ka1; RSK 1; RSK 1 p90; RSK; RSK-1; RSK1; S6K alpha 1; S6K-alpha-1; anti-RSK1p90 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-RSK1 p90 (Thr348) Antibody detects endogenous levels of RSK1 p90 only when phosphorylated at Thr348.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-RSK1p90 antibody	IHC (Immunohistochemistry), ELISA (Peptide ELISA)
Application Notes	IHC: 1:50-1:200 Peptide ELISA: 1:20,000-1:40,000
Immunogen	A synthesized peptide derived from human RSK1 p90 around the phosphorylation site of Thr348.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	Activated by phosphorylation at Ser-221 by PDPK1. Autophosphorylated on Ser-380, as part of the activation process. May be phosphorylated at Thr-359 and Ser-363 by MAPK1/ERK2 and MAPK3/ERK1. N-terminal myristoylation results in an activated kinase in the absence of added growth factors.
Subunit Structure	Forms a complex with either MAPK1/ERK2 or MAPK3/ERK1 in quiescent cells. Transiently dissociates following mitogenic stimulation. Interacts with ETV1/ER81 and FGFR1.
Similarity	Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.
Subcellular Location	Nucleus. Cytoplasm.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IHC (Immunohistochemistry)

At 1/100 staining Human kidney cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)

At 1/100 staining Human kidney cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)

At 1/100 staining Human ovarian cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)

At 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)

Staining Rat kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)

At 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-RSK1p90 antibody	Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1) signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro-apoptotic function of BAD and DAPK1. In fibroblast, is required for EGF-stimulated phosphorylation of CREB1, which results in the subsequent transcriptional activation of several immediate-early genes. In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP. Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity. Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the pre-initiation complex. In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation. Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin-sensitive signaling independently of the PI3K/AKT pathway. Mediates cell survival by phosphorylating the pro-apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function. Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4). Mediates induction of hepatocyte proliferation by TGFA through phosphorylation of CEBPB (By similarity). Is involved in cell cycle regulation by phosphorylating the CDK inhibitor CDKN1B, which promotes CDKN1B association with 14-3-3 proteins and prevents its translocation to the nucleus and inhibition of G1 progression. Phosphorylates EPHA2 at 'Ser-897', the RPS6KA-EPHA2 signaling pathway controls cell migration.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	55743134
NCBI GeneID	6195
NCBI Accession #	NP_001006666.1
NCBI GenBank Nucleotide #	NM_001006665.1
UniProt Accession #	Q15418; Q5SVM5; Q5SVM8; Q5SVM9; Q96C05; Q9BQK2; A6NGG4; A8K9K7; B2RDY8; B7Z5J0
Molecular Weight	82,723 Da
NCBI Official Full Name	ribosomal protein S6 kinase alpha-1 isoform b
NCBI Official Synonym Full Names	ribosomal protein S6 kinase, 90kDa, polypeptide 1
NCBI Official Symbol	RPS6KA1
NCBI Official Synonym Symbols	RSK; HU-1; RSK1; MAPKAPK1A
NCBI Protein Information	ribosomal protein S6 kinase alpha-1; RSK-1; p90S6K; p90RSK1; p90-RSK 1; MAPKAPK-1a; S6K-alpha 1; S6K-alpha-1; MAPKAP kinase 1a; ribosomal S6 kinase 1; MAPK-activated protein kinase 1a; ribosomal protein S6 kinase alpha 1; 90 kDa ribosomal protein S6 kinas
UniProt Protein Name	Ribosomal protein S6 kinase alpha-1
UniProt Gene Name	RPS6KA1
UniProt Synonym Gene Names	MAPKAPK1A; RSK1; S6K-alpha-1; p90-RSK 1; p90RSK1; p90S6K; MAPK-activated protein kinase 1a; MAPKAP kinase 1a; MAPKAPK-1a; RSK-1
UniProt Entry Name	KS6A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rsk1-p90-antibody-immunohistochemistry-peptide-elisa-elisa/31316>