

Rabbit anti-Human, Mouse MMP2 Polyclonal Antibody | anti-MMP2 antibody

MMP2 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA313243	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MMP2; Polyclonal Antibody; MMP2 Rabbit Polyclonal Antibody; CLG4; CLG4A; MMP-2; MMP-II; MONA; TBE-1; anti-MMP2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-MMP2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human MMP2 (NP_004521.1).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using MMP2 at dilution of 1:100. Blue: DAPI for nuclear staining.


WB (Western Blot)

Western blot analysis of extracts of various cell lines, using MMP2 at 1:3000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP2 antibody	This gene is a member of the matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of cleaving components of the extracellular matrix and molecules involved in signal transduction. The protein encoded by this gene is a gelatinase A, type IV collagenase, that contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or, intracellularly by its S-glutathiolation with no requirement for proteolytical removal of the pro-domain. This protein is thought to be involved in multiple pathways including roles in the nervous system, endometrial menstrual breakdown, regulation of vascularization, and metastasis. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms.
Product Categories/Family for anti-MMP2 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	189217853
NCBI GeneID	4313
NCBI Accession #	NP_001121363.1
NCBI GenBank Nucleotide #	NM_001127891.1
UniProt Accession #	P08253; Q9UCJ8; B2R6U1; B4DWH3; E9PE45
Molecular Weight	73,882 Da
NCBI Official Full Name	72 kDa type IV collagenase isoform b
NCBI Official Synonym Full Names	matrix metallopeptidase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase)
NCBI Official Symbol	MMP2
NCBI Official Synonym Symbols	CLG4; MONA; CLG4A; TBE-1; MMP-II
NCBI Protein Information	72 kDa type IV collagenase; MMP-2; gelatinase A; 72 kDa gelatinase; collagenase type IV-A; neutrophil gelatinase; matrix metalloproteinase-2; matrix metalloproteinase-II
UniProt Protein Name	72 kDa type IV collagenase
UniProt Gene Name	MMP2
UniProt Synonym Gene Names	CLG4A; MMP-2
UniProt Entry Name	MMP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.