

Rabbit FLT3 Polyclonal Antibody | anti-FLT3 antibody

FLT3 Rabbit Polyclonal Antibody

CATALOG # AAA313300	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

FLT3; Polyclonal Antibody; FLT3 Rabbit Polyclonal Antibody; CD135; FLK-2; FLK2; STK1; anti-FLT3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-FLT3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human FLT3 (NP_004110.2).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver cancer using FLT3 at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat pancreas using FLT3 at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using FLT3 at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-FLT3 antibody	This gene encodes a class III receptor tyrosine kinase that regulates hematopoiesis. This receptor is activated by binding of the fms-related tyrosine kinase 3 ligand to the extracellular domain, which induces homodimer formation in the plasma membrane leading to autophosphorylation of the receptor. The activated receptor kinase subsequently phosphorylates and activates multiple cytoplasmic effector molecules in pathways involved in apoptosis, proliferation, and differentiation of hematopoietic cells in bone marrow. Mutations that result in the constitutive activation of this receptor result in acute myeloid leukemia and acute lymphoblastic leukemia.
Product Categories/Family for anti-FLT3 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	121114304
NCBI GeneID	2322
NCBI Accession #	NP_004110.2
NCBI GenBank Nucleotide #	NM_004119.2
UniProt Accession #	P36888; Q13414; A0AVG9; B7ZLT7; B7ZLT8; F5H0A0
Molecular Weight	112,903 Da
NCBI Official Full Name	receptor-type tyrosine-protein kinase FLT3
NCBI Official Synonym Full Names	fms-related tyrosine kinase 3
NCBI Official Symbol	FLT3
NCBI Official Synonym Symbols	FLK2; STK1; CD135; FLK-2
NCBI Protein Information	receptor-type tyrosine-protein kinase FLT3; STK-1; CD135 antigen; FL cytokine receptor; fetal liver kinase 2; fms-like tyrosine kinase 3; stem cell tyrosine kinase 1; growth factor receptor tyrosine kinase type III
UniProt Protein Name	Receptor-type tyrosine-protein kinase FLT3
UniProt Gene Name	FLT3
UniProt Synonym Gene Names	CD135; FLK2; STK1; FLK-2; FLT-3; STK-1
UniProt Entry Name	FLT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/flt3-antibody-immunohistochemistry-western-blot-elisa/313300>