

Rabbit CD68 Polyclonal Antibody | anti-CD68 antibody

CD68 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA313388	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD68; Polyclonal Antibody; CD68 Rabbit Polyclonal Antibody; GP110; LAMP4; SCARD1; anti-CD68 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-CD68 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human CD68 (NP_001242.2).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




WB (Western Blot)

Western blot analysis of extracts of Rat thymus, using CD68 Polyclonal at 1:1000 dilution.




WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CD68 Polyclonal at 1:1000 dilution.



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Mouse spleen using CD68 at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human oophoroma using CD68 at dilution of 1:100 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-CD68 antibody	This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms.
Product Categories/Family for anti-CD68 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	298665
NCBI GeneID	968
UniProt Accession #	P34810 ; Q53HR6 ; Q53XI3 ; Q96BI7 ; B4DVT4
Molecular Weight	37,408 Da
NCBI Official Full Name	CD68
NCBI Official Synonym Full Names	CD68 molecule
NCBI Official Symbol	CD68
NCBI Official Synonym Symbols	GP110; LAMP4; SCARD1
NCBI Protein Information	macrosialin; CD68 antigen; macrophage antigen CD68; scavenger receptor class D, member 1
UniProt Protein Name	Macrosialin
UniProt Gene Name	CD68
UniProt Entry Name	CD68_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd68-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/313388>