

Rabbit WT1 Polyclonal Antibody | anti-WT1 antibody

WT1 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA313650	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

WT1; Polyclonal Antibody; WT1 Rabbit Polyclonal Antibody; AWT1; EWS-WT1; GUD; NPHS4; WAGR; WIT-2; WT33; anti-WT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-WT1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human WT1
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil using WT1 at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat testis using WT1 at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using WT1 at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-WT1 antibody

This gene encodes a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilms tumor. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation codon upstream of, and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-restricted and developmentally regulated.

Product Categories/Family for anti-WT1 antibody

[Primary Antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	65507817
NCBI GeneID	7490
NCBI Accession #	NP_077742.2
NCBI GenBank Nucleotide #	NM_024424.3
UniProt Accession #	P19544 ; Q15881 ; Q16256 ; Q16575 ; Q4VXV4 ; Q4VXV5 ; Q4VXV6 ; Q8IYZ5 ; A8K6S1 ; B3KSA5
Molecular Weight	49,188 Da
NCBI Official Full Name	Wilms tumor protein isoform B
NCBI Official Synonym Full Names	Wilms tumor 1
NCBI Official Symbol	WT1
NCBI Official Synonym Symbols	GUD; AWT1; WAGR; WT33; NPHS4; WIT-2; EWS-WT1
NCBI Protein Information	Wilms tumor protein; amino-terminal domain of EWS last three zinc fingers of the DNA-binding domain of WT1
UniProt Protein Name	Wilms tumor protein
UniProt Gene Name	WT1
UniProt Entry Name	WT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.