

Rabbit CSK Polyclonal Antibody | anti-CSK antibody

Phospho-CSK (Tyr304) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31403	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA


PRODUCT SYNONYMS

CSK; Polyclonal Antibody; Phospho-CSK (Tyr304) Antibody; C SRC; C SRC kinase; C src Tyrosine Kinase; C-SRC kinase; c-src tyrosine kinase; Csk A; CSK_HUMAN; CYTOPLASMIC TYROSINE KINASE; EC 2.7.10.2; MGC112926; MGC117393; MGC154049; P60 Src; Protein tyrosine kinase CYL; Protein-tyrosine kinase CYL; Proto oncogene tyrosine protein kinase; Tyrosine protein kinase CSK; Tyrosine-protein kinase CSK; zgc:154049; anti-CSK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-CSK (Tyr304) Antibody detects endogenous levels of CSK only when phosphorylated at Tyr304. Tissue Specificity: Expressed in lung and macrophages.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Applicable Applications for anti-CSK antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA (Peptide ELISA)
Application Notes	WB: 1:1000-3000 IHC: 1:50-1:200 IF/ICC: 1:100-1:500 Peptide ELISA: 1:20,000-1:40,000 The optimal dilutions should be determined by the end user. IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.
Immunogen	A synthesized peptide derived from human CSK around the phosphorylation site of Tyr304.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



Application Data

At 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Red), diluted at 1/600, was used as secondary antibody.



IHC (Immunohistochemistry)

At 1/200 staining Human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)

At 1/200 staining Mouse heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)

At 1/200 staining Rat spleen tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)

At 1/200 staining Human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)

At 1/200 staining Mouse kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.

NCBI AND UNIPROT INFORMATION

NCBI GI #	187475373
NCBI GeneID	1445
NCBI Accession #	NP_001120662.1
NCBI GenBank Nucleotide #	NM_001127190.1
UniProt Accession #	P41240 ; Q2M3N2 ; Q6FGZ6
Molecular Weight	55KD
NCBI Official Full Name	tyrosine-protein kinase CSK
NCBI Official Synonym Full Names	c-src tyrosine kinase
NCBI Official Symbol	CSK
NCBI Protein Information	tyrosine-protein kinase CSK; C-Src kinase; protein-tyrosine kinase CYL
UniProt Protein Name	Tyrosine-protein kinase CSK
UniProt Gene Name	CSK
UniProt Entry Name	CSK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/csk-antibody-western-blot-immunohistochemistry-immunofluorescence-immunocytochemistry-peptide-elisa-elisa/31403>