

anti-Human CD63 Monoclonal Antibody | anti-CD63 antibody

CD63 Rabbit mAb

CATALOG # AAA314328	HOST Recombinant Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------------------	----------------------------	---

PRODUCT SYNONYMS

CD63; Monoclonal Antibody; CD63 Rabbit mAb; MLA1; ME491; LAMP-3; OMA81H; TSPAN30; anti-CD63 antibody

PRODUCT SPECIFICATIONS

Host	Recombinant Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	S08-7C9
Purity/Purification	Affinity Purified
Form/Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration	0.3mg/ml (varies by lot)
Applicable Applications for anti-CD63 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human CD63
Conjugation	Unconjugated
Modification	Unmodification
Preparation and Storage	Store at 4 degree C short term. Aliquot and store at -20 degree C long term. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of CD63 in paraffin-embedded Human breast cancer tissue using CD63 Rabbit mAb at dilution 1/100



WB (Western Blot)

Western blot detection of CD63 in HeLa cell lysates using CD63 Rabbit mAb(1:1000 diluted). Predicted band size: 26kDa. Observed band size: 26-65kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD63 antibody	Functions as cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2. Plays a role in VEGFA signaling via its role in regulating the internalization of KDR/VEGFR2. Plays a role in intracellular vesicular transport processes, and is required for normal trafficking of the PMEL luminal domain that is essential for the development and maturation of melanocytes. Plays a role in the adhesion of leukocytes onto endothelial cells via its role in the regulation of SELP trafficking. May play a role in mast cell degranulation in response to Ms4a2/FceRI stimulation, but not in mast cell degranulation in response to other stimuli.
Product Categories/Family for anti-CD63 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	189384
NCBI GeneID	967
UniProt Accession #	P08962; Q5TZP3; Q8N6Z9; Q9UCG6; F8VZE2
Molecular Weight	23,430 Da
NCBI Official Full Name	melanoma-associated antigen
NCBI Official Synonym Full Names	CD63 molecule
NCBI Official Symbol	CD63
NCBI Official Synonym Symbols	MLA1; ME491; LAMP-3; OMA81H; TSPAN30
NCBI Protein Information	CD63 antigen
UniProt Protein Name	CD63 antigen
UniProt Gene Name	CD63
UniProt Synonym Gene Names	MLA1; TSPAN30; LAMP-3; Tspan-30

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.