

NF kappa B p100 Monoclonal Antibody | anti-NFkappaB antibody

NF kappa B p100 Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA314422	Recombinant Rabbit	Human, Mouse, Rat	Immunofluorescence, Western Blot


PRODUCT SYNONYMS

NF kappa B p100; Monoclonal Antibody; NF kappa B p100 Rabbit mAb; p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100; anti-NFkappaB antibody

PRODUCT SPECIFICATIONS


Host	Recombinant Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	S04-4D6
Purity/Purification	Affinity Purified
Form/Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration	0.3mg/ml (varies by lot)
Applicable Applications for anti-NFkappaB antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant protein of human NFkB p100
Conjugation	Unconjugated
Modification	Unmodification
Preparation and Storage	Store at 4 degree C short term. Aliquot and store at -20 degree C long term. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence of NFkB p100(green) in HeLa cells using NFkB p100 Rabbit mAb at dilution 1/200, and DAPI(blue)



WB (Western Blot)

Western blot detection of NFkB p100/NFKB2 in C6,3T3,HeLa cell lysates using NFkB p100/NFKB2 Rabbit mAb(1:1000 diluted).Predicted band size:97kDa.Observed band size:100kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-NFkappaB antibody	NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. In a non-canonical activation pathway, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. The NF-kappa-B heterodimeric RelB-p52 complex is a transcriptional activator. The NF-kappa-B p52-p52 homodimer is a transcriptional repressor. NFKB2 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p100 and generation of p52 by a cotranslational processing. The proteasome-mediated process ensures the production of both p52 and p100 and preserves their independent function. p52 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. p52 and p100 are respectively the minor and major form; the processing of p100 being relatively poor. Isoform p49 is a subunit of the NF-kappa-B protein complex, which stimulates the HIV enhancer in synergy with p65. In concert with RELB, regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer.
Product Categories/Family for anti-NFkappaB antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	117320531
NCBI GeneID	4791
NCBI Accession #	NP_001070962.1
NCBI GenBank Nucleotide #	NM_001077494.2
UniProt Accession #	Q00653; Q04860; Q9BU75; Q9H471; Q9H472; A8K9D9; D3DR83; AAB23437
Molecular Weight	96,749 Da
NCBI Official Full Name	nuclear factor NF-kappa-B p100 subunit isoform a
NCBI Official Synonym Full Names	nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)
NCBI Official Symbol	NFKB2
NCBI Official Synonym Symbols	p52; p105; H2TF1; LYT10; LYT-10; NF-kB2
NCBI Protein Information	nuclear factor NF-kappa-B p100 subunit; oncogene Lym-10; DNA-binding factor KBF2; lymphocyte translocation chromosome 10 protein; nuclear factor of kappa light chain gene enhancer in B-cells 2
UniProt Protein Name	Nuclear factor NF-kappa-B p100 subunit
UniProt Gene Name	NFKB2
UniProt Synonym Gene Names	LYT10; Lym10
UniProt Entry Name	NFKB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.