

PKC delta Monoclonal Antibody | anti-PKCdelta antibody

PKC delta Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA314436	Recombinant Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PKC delta; Monoclonal Antibody; PKC delta Rabbit mAb; Pkcd; PKC[d]; A1385711; PKCdelta; D14Ert420e; anti-PKCdelta antibody

PRODUCT SPECIFICATIONS


Host	Recombinant Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	S09-8H6
Purity/Purification	Affinity Purified
Form/Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration	0.3mg/ml (varies by lot)
Applicable Applications for anti-PKCdelta antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human PKC delta
Conjugation	Unconjugated
Modification	Unmodification
Preparation and Storage	Store at 4 degree C short term. Aliquot and store at -20 degree C long term. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of PKC delta in paraffin-embedded Human colon cancer tissue using PKC delta Rabbit mAb at dilution 1/100



WB (Western Blot)

Western blot detection of PKC delta in K562,C6,3T3 cell lysates using PKC delta Rabbit mAb(1:1000 diluted).Predicted band size:78kDa.Observed band size:78kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PKCdelta antibody	Calcium-independent, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase that plays contrasting roles in cell death and cell survival by functioning as a pro-apoptotic protein during DNA damage-induced apoptosis, but acting as an anti-apoptotic protein during cytokine receptor-initiated cell death, is involved in tumor suppression, is required for oxygen radical production by NADPH oxidase and acts as positive or negative regulator in platelet functional responses. Negatively regulates B cell proliferation and also has an important function in self-antigen induced B cell tolerance induction. Upon DNA damage, activates the promoter of the death-promoting transcription factor BCLAF1/Btf to trigger BCLAF1-mediated p53/TP53 gene transcription and apoptosis. In response to oxidative stress, interact with and activate CHUK/IKKA in the nucleus, causing the phosphorylation of p53/TP53. In the case of ER stress or DNA damage-induced apoptosis, can form a complex with the tyrosine-protein kinase ABL1 which trigger apoptosis independently of p53/TP53. In cytosol can trigger apoptosis by activating MAPK11 or MAPK14, inhibiting AKT1 and decreasing the level of X-linked inhibitor of apoptosis protein (XIAP), whereas in nucleus induces apoptosis via the activation of MAPK8 or MAPK9. Upon ionizing radiation treatment, is required for the activation of the apoptosis regulators BAX and BAK, which trigger the mitochondrial cell death pathway. Can phosphorylate MCL1 and target it for degradation which is sufficient to trigger for BAX activation and apoptosis. Is required for the control of cell cycle progression both at G1/S and G2/M phases. Mediates phorbol 12-myristate 13-acetate (PMA)-induced inhibition of cell cycle progression at G1/S phase by up-regulating the CDK inhibitor CDKN1A/p21 and inhibiting the cyclin CCNA2 promoter activity. In response to UV irradiation can phosphorylate CDK1, which is important for the G2/M DNA damage checkpoint activation. Can protect glioma cells from the apoptosis induced by TNFSF10/TRAIL, probably by inducing increased phosphorylation and subsequent activation of AKT1. Can also act as tumor suppressor upon mitogenic stimulation with PMA or TPA. In N-formyl-methionyl-leucyl-phenylalanine (fMLP)-treated cells, is required for NCF1 (p47-phox) phosphorylation and activation of NADPH oxidase activity, and regulates TNF-elicited superoxide anion production in neutrophils, by direct phosphorylation and activation of NCF1 or indirectly through MAPK1/3 (ERK1/2) signaling pathways. May also play a role in the regulation of NADPH oxidase activity in eosinophil after stimulation with IL5, leukotriene B4 or PMA. In collagen-induced platelet aggregation, acts a negative regulator of filopodia formation and actin polymerization by interacting with and negatively regulating VASP phosphorylation. Downstream of PAR1, PAR4 and CD36/GP4 receptors, regulates differentially platelet dense granule secretion; acts as a positive regulator in PAR-mediated granule secretion, whereas it negatively regulates CD36/GP4-mediated granule release. Phosphorylates MUC1 in the C-terminal and regulates the interaction between MUC1 and beta-catenin. The catalytic subunit phosphorylates 14-3-3 proteins (YWHAB, YWHAZ and YWHAH) in a sphingosine-dependent fashion. Phosphorylates ELAVL1 in response to angiotensin-2 treatment .
Product Categories/Family for anti-PKCdelta antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	6755082
NCBI GeneID	18753
NCBI Accession #	NP_035233.1
NCBI GenBank Nucleotide #	NM_011103.3
UniProt Accession #	P28867; Q91V85; Q9Z333
Molecular Weight	80,294 Da
NCBI Official Full Name	protein kinase C delta type
NCBI Official Synonym Full Names	protein kinase C, delta
NCBI Official Symbol	Prkcd
NCBI Official Synonym Symbols	Pkcd; PKC[d]; AI385711; PKCdelta; D14Ert420e
NCBI Protein Information	protein kinase C delta type; nPKC-delta; protein kinase C delta variant IX; protein kinase C, delta IV; protein kinase C, delta V; protein kinase C[d]; tyrosine-protein kinase PRKCD
UniProt Protein Name	Protein kinase C delta type
UniProt Gene Name	Prkcd

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/pkc-delta-antibody-clone-s09-8h6-immunohistochemistry-western-blot-elisa/314436>