

Rabbit anti-Human CD227/MUC1 Polyclonal Antibody | anti-MUC1 antibody

CD227/MUC1 (Phospho-Tyr1229) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA316434	Rabbit	Human	ELISA, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD227/MUC1; Polyclonal Antibody; CD227/MUC1 (Phospho-Tyr1229) Antibody; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; CD227; EMA; Episialin; H23AG; MUC-1; mucin 1; Mucin 1 precursor; transmembrane; Peanut-reactive urinary mucin; PEM; PEMT; Polymorphic epithelial mucin; PUM; Tumor-associated epi; anti-MUC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	CD227/MUC1 (Phospho-Tyr1229) Antibody detects endogenous levels of CD227/MUC1 only when phosphorylated at Tyr1229.
Purity/Purification	Purified from rabbit antiserum by affinity-chromatography using phospho peptide. The antibody against non-phospho peptide was removed by chromatography using corresponding non-phospho peptide.
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1 mg/ml (varies by lot)
Sequence Length	259
Applicable Applications for anti-MUC1 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Target Modification	Phospho
Modification Sites	H:Y1229
Immunogen	The antiserum was produced against synthesized peptide derived from human CD227/MUC1 around the phosphorylation site of Tyr1229.
Immunogen Range	1201-1250
Preparation and Storage	Stable at -20 degree C for at least 1 year.

IMAGES




WB (Western Blot)

Western blot analysis of lysates from HepG2 cells treated with PMA 125ng/ml 30', using CD227/MUC1 (Phospho-Tyr1229) Antibody. The lane on the right is blocked with the phospho peptide.



IF (Immunofluorescence)

Immunofluorescence analysis of HepG2 cells, using CD227/MUC1 (Phospho-Tyr1229) Antibody. The picture on the right is blocked with the phospho peptide.



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using CD227/MUC1 (Phospho-Tyr1229) Antibody. The picture on the right is blocked with the phospho peptide.

NCBI AND UNIPROT INFORMATION

NCBI GI #	67189007
NCBI GeneID	4582
NCBI Accession #	NP_001018016.1
NCBI GenBank Nucleotide #	NM_001018016.2
UniProt Accession #	P15941; P13931; P15942; P17626; A5YRV1; A6ZID9; A6ZIE0; B1AVQ8; B1AVR0; B6ECA1; E7ESE5; E7EUG9
Molecular Weight	122 kDa
NCBI Official Full Name	mucin-1 isoform 2
NCBI Official Synonym Full Names	mucin 1, cell surface associated
NCBI Official Symbol	MUC1
NCBI Official Synonym Symbols	EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC
NCBI Protein Information	mucin-1
UniProt Protein Name	Mucin-1
UniProt Gene Name	MUC1
UniProt Synonym Gene Names	PUM; MUC-1; CA 15-3; KL-6; PUM; PEM; EMA; MUC1-NT; MUC1-alpha; MUC1-beta

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

