

Rabbit CD66a/b/c Polyclonal Antibody | anti-CD66a/b/c antibody

CD66a/b/c Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318796	Rabbit	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

CD66a/b/c; Polyclonal Antibody; CD66a/b/c Rabbit Polyclonal Antibody; BH6842; Carcinoembryonic antigen-related cell adhesion molecule 1/6/8 (CD antigen CD66a/b/c); anti-CD66a/b/c antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous CD66a/b/c
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-CD66a/b/c antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 130-190
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human-kidney, antibody was diluted at 1:100



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human-stomach, antibody was diluted at 1:100



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded Human-stomach, antibody was diluted at 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-CD66a/b/c antibody	This gene encodes a member of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily. Two subgroups of the CEA family, the CEA cell adhesion molecules and the pregnancy-specific glycoproteins, are located within a 1.2 Mb cluster on the long arm of chromosome 19. Eleven pseudogenes of the CEA cell adhesion molecule subgroup are also found in the cluster. The encoded protein was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. Multiple cellular activities have been attributed to the encoded protein, including roles in the differentiation and arrangement of tissue thre
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	399116
NCBI GeneID	634
UniProt Accession #	P13688; P40199; P31997
Molecular Weight	526
NCBI Official Full Name	Carcinoembryonic antigen-related cell adhesion molecule 1
NCBI Official Synonym Full Names	carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein)
NCBI Official Symbol	CEACAM1
NCBI Official Synonym Symbols	BGP; BGP1; BGPI
NCBI Protein Information	carcinoembryonic antigen-related cell adhesion molecule 1; antigen CD66; CD66a antigen
UniProt Protein Name	Carcinoembryonic antigen-related cell adhesion molecule 1
UniProt Gene Name	CEACAM1
UniProt Synonym Gene Names	BGP; BGP1; BGP-1
UniProt Entry Name	CEAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.