

Rabbit FAS-L Polyclonal Antibody | anti-FAS-L antibody

FAS-L Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318817	Rabbit	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

FAS-L; Polyclonal Antibody; FAS-L Rabbit Polyclonal Antibody; Tumor necrosis factor ligand superfamily member 6 (Apoptosis antigen ligand)(APTL)(CD95 ligand)(CD95-L)(Fas antigen ligand)(Fas ligand)(FasL)(CD antigen CD178) [Cleaved into: Tumor necrosis factor ligand superfamily member 6; membrane form; Tumor necrosis factor ligand superfamily member 6; soluble form (Receptor-binding FasL ectodomain)(Soluble Fas ligand)(sFasL); ADAM10-processed FasL form (APL); FasL intracellular domain (FasL ICD)(SPPL2A-processed FasL form)(SPA)]; BH6842; anti-FAS-L antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous FAS-L
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-FAS-L antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 121-170
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded Human-colon-cancer, antibody was diluted at 1:100



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded Human-colon-cancer, antibody was diluted at 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-FAS-L antibody	This gene is a member of the tumor necrosis factor superfamily. The primary function of the encoded transmembrane protein is the induction of apoptosis triggered by binding to FAS. The FAS/FASLG signaling pathway is essential for immune system regulation, including activation-induced cell death (AICD) of T cells and cytotoxic T lymphocyte induced cell death. It has also been implicated in the progression of several cancers. Defects in this gene may be related to some cases of systemic lupus erythematosus (SLE). Alternatively spliced transcript variants have been described. [provided by RefSeq, Nov 2014]
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557329
NCBI GeneID	356
NCBI Accession #	NP_000630.1
NCBI GenBank Nucleotide #	NM_000639.1
UniProt Accession #	P48023; Q9BZP9
Molecular Weight	31,485 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 6
NCBI Official Synonym Full Names	Fas ligand (TNF superfamily, member 6)
NCBI Official Symbol	FASLG
NCBI Official Synonym Symbols	APTL; FASL; CD178; CD95L; ALPS1B; CD95-L; TNFSF6; APT1LG1
NCBI Protein Information	tumor necrosis factor ligand superfamily member 6; CD95 ligand; fas antigen ligand; apoptosis antigen ligand; apoptosis (APO-1) antigen ligand 1; tumor necrosis factor (ligand) superfamily, member 6
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 6
UniProt Gene Name	FASLG
UniProt Synonym Gene Names	APT1LG1; CD95L; FASL; TNFSF6; APTL; CD95-L; Fas ligand; FasL; sFasL; APL; FasL ICD; SPA
UniProt Entry Name	TNFL6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fas-l-antibody-immunofluorescence-immunohistochemistry-elisa/318817>