

Rabbit Inhibin beta-B Polyclonal Antibody | anti-INHBB antibody

Inhibin beta-B Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318830	Rabbit	Human, Mouse, Rat	ELISA, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Inhibin beta-B; Polyclonal Antibody; Inhibin beta-B Rabbit Polyclonal Antibody; INHBB; Inhibin beta B chain; Activin beta-B chain; BH6842; anti-INHBB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Inhibin-beta-B Polyclonal Antibody detects endogenous levels of Inhibin-beta-B protein.
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-INHBB antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthesized peptide derived from Inhibin beta B chain at AA range: 351-400
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




WB (Western Blot)

Western Blot analysis of HeLa cells using Inhibin beta-B Polyclonal Antibody. Antibody was diluted at 1:2000. Secondary antibody was diluted at 1:20000




WB (Western Blot)

Western blot analysis of YT5408wb4147442094 lysis using INHBB Antibody antibody. Antibody was diluted at 1:2000. Secondary antibody was diluted at 1:20000




IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-INHBB antibody

This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily of proteins. The encoded preproprotein is proteolytically processed to generate a subunit of the dimeric activin and inhibin Protein complexes. These complexes activate and inhibit, respectively, follicle stimulating hormone secretion from the pituitary gland. Polymorphisms near this gene are associated with pre-eclampsia in female human patients. [provided by RefSeq, Aug 2016]

NCBI AND UNIPROT INFORMATION

NCBI GI #	154813204
NCBI GeneID	3625
NCBI Accession #	NP_002184.2
NCBI GenBank Nucleotide #	NM_002193.2
UniProt Accession #	P09529 ; Q53T31 ; Q8N1D3
Molecular Weight	45,122 Da
NCBI Official Full Name	inhibin beta B chain preproprotein
NCBI Official Synonym Full Names	inhibin, beta B
NCBI Official Symbol	INHBB
NCBI Protein Information	inhibin beta B chain; Inhibin, beta-2; activin beta-B chain; activin AB beta polypeptide
UniProt Protein Name	Inhibin beta B chain
UniProt Gene Name	INHBB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/inhibin-beta-b-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/318830>