

Rabbit anti-Human, Mouse MIP-3alpha Polyclonal Antibody | anti-MIP-3alpha antibody

MIP-3alpha Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318838	Rabbit	Human, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

MIP-3alpha; Polyclonal Antibody; MIP-3alpha Rabbit Polyclonal Antibody; C-C motif chemokine 20 (Beta-chemokine exodus-1)(CC chemokine LARC)(Liver and activation-regulated chemokine)(Macrophage inflammatory protein 3 alpha)(MIP-3alphaalpha)(Small-inducible cytokine A20) [Cleaved into: CCL20 (1-67); CCL20 (1-64); CCL20 (2-70)]; BH6842; anti-MIP-3alpha antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous MIP-3alpha
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-MIP-3alpha antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 31-80
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES





IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:200

IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:200

ADDITIONAL INFORMATION

Related Product Information for anti-MIP-3alpha antibody

This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene displays chemotactic activity for lymphocytes and can repress proliferation of myeloid progenitors. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]

NCBI AND UNIPROT INFORMATION

NCBI GI #	2829668
NCBI GeneID	6364
UniProt Accession #	P78556; Q53S51; Q99664
Molecular Weight	8025.5
NCBI Official Full Name	C-C motif chemokine 20
NCBI Official Synonym Full Names	C-C motif chemokine ligand 20
NCBI Official Symbol	CCL20
NCBI Official Synonym Symbols	CKb4; LARC; ST38; MIP3A; Exodus; MIP-3a; SCYA20; MIP-3-alpha
NCBI Protein Information	C-C motif chemokine 20
UniProt Protein Name	C-C motif chemokine 20
UniProt Gene Name	CCL20
UniProt Synonym Gene Names	LARC; MIP3A; SCYA20; MIP-3-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mip-3alpha-antibody-immunofluorescence-immunohistochemistry-elisa/318838>