

Rabbit ENA-78 Polyclonal Antibody | anti-ENA-78 antibody

ENA-78 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318845	Rabbit	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

ENA-78; Polyclonal Antibody; ENA-78 Rabbit Polyclonal Antibody; C-X-C motif chemokine 5 (ENA-78 (1-78))(Epithelial-derived neutrophil-activating protein 78)(Neutrophil-activating peptide ENA-78)(Small-inducible cytokine B5) [Cleaved into: ENA-78 (8-78); ENA-78 (9-78)]; BH6842; anti-ENA-78 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous ENA-78
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-ENA-78 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 65-114
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-pancreas-cancer, antibody was diluted at 1:200



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-pancreas-cancer, antibody was diluted at 1:200

ADDITIONAL INFORMATION

Related Product Information for anti-ENA-78 antibody	This gene encodes a protein that is a member of the CXC subfamily of chemokines. Chemokines, which recruit and activate leukocytes, are classified by function (inflammatory or homeostatic) or by structure. This protein is proposed to bind the G-protein coupled receptor chemokine (C-X-C motif) receptor 2 to recruit neutrophils, to promote angiogenesis and to remodel connective tissues. This protein is thought to play a role in cancer cell proliferation, migration, and invasion. [provided by RefSeq, May 2013]
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	471243
NCBI GeneID	6374
UniProt Accession #	P42830; Q96QE1
Molecular Weight	11,972 Da
NCBI Official Full Name	ENA-78
NCBI Official Synonym Full Names	chemokine (C-X-C motif) ligand 5
NCBI Official Symbol	CXCL5
NCBI Official Synonym Symbols	SCYB5; ENA-78
NCBI Protein Information	C-X-C motif chemokine 5; neutrophil-activating protein 78; neutrophil-activating peptide ENA-78; epithelial-derived neutrophil-activating protein 78; small inducible cytokine subfamily B (Cys-X-Cys), member 5 (epithelial-derived neutrophil-activating pept
UniProt Protein Name	C-X-C motif chemokine 5
UniProt Gene Name	CXCL5
UniProt Synonym Gene Names	ENA78; SCYB5
UniProt Entry Name	CXCL5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ena-78-antibody-immunofluorescence-immunohistochemistry-elisa/318845>