

Rabbit Bob 1 Polyclonal Antibody | anti-Bob1 antibody

Bob 1 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318866	Rabbit	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

Bob 1; Polyclonal Antibody; Bob 1 Rabbit Polyclonal Antibody; POU domain class 2-associating factor 1 (B-cell-specific coactivator OBF-1)(BOB-1)(OCA-B)(OCT-binding factor 1); BH6842; anti-Bob1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous Bob 1
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-Bob1 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 201-250
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES





IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200




IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4 degree overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



IHC (Immunohistochemistry)
 Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4 degree overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

IHC (Immunohistochemistry)
 Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4 degree overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



IHC (Immunohistochemistry)
 Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4 degree overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

NCBI AND UNIPROT INFORMATION

NCBI GI #	167900478
NCBI GeneID	5450
NCBI Accession #	NP_006226.2
NCBI GenBank Nucleotide #	NM_006235.2
UniProt Accession #	Q16633; Q14983; B2R8Z9; AAB34937
Molecular Weight	27,436 Da
NCBI Official Full Name	POU domain class 2-associating factor 1
NCBI Official Synonym Full Names	POU class 2 associating factor 1
NCBI Official Symbol	POU2AF1
NCBI Official Synonym Symbols	BOB1; OBF1; OCAB; OBF-1
NCBI Protein Information	POU domain class 2-associating factor 1; B-cell-specific coactivator OBF-1; BOB-1; OCA-B; OCT-binding factor 1; POU domain, class 2, associating factor 1
UniProt Protein Name	POU domain class 2-associating factor 1
UniProt Gene Name	POU2AF1
UniProt Synonym Gene Names	OBF1
UniProt Entry Name	OBF1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/bob-1-antibody-immunofluorescence-immunohistochemistry-elisa/318866>