

Rabbit LIMK-2 Polyclonal Antibody | anti-LIMK-2 antibody

LIMK-2 (Phospho-Thr505) Antibody

CATALOG # AAA319461	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

LIMK-2; Polyclonal Antibody; LIMK-2 (Phospho-Thr505) Antibody; LIM domain kinase 2; LIMK2; anti-LIMK-2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-LIMK-2 (T505) Polyclonal Antibody detects endogenous levels of LIMK-2 protein only when phosphorylated at T505.
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-LIMK-2 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthesized phospho-peptide around the phosphorylation site of human LIMK-2 (phospho Thr505)
Modification	Phospho Specific
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



WB (Western Blot)

Western Blot analysis of various cells using Phospho-LIMK-2 (T505) Polyclonal Antibody



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4 degree overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

ADDITIONAL INFORMATION

Related Product Information for anti-LIMK-2 antibody	There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Product Categories/Family for anti-LIMK-2 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	73390140
NCBI GeneID	3985
NCBI Accession #	NP_001026971.1
NCBI GenBank Nucleotide #	NM_001031801.1
UniProt Accession #	P53671; Q7KZ80; Q7L3H5; Q96E10; Q99464; Q9UFU0; A8K6H5
Molecular Weight	72,232 Da
NCBI Official Full Name	LIM domain kinase 2 isoform 1
NCBI Official Synonym Full Names	LIM domain kinase 2
NCBI Official Symbol	LIMK2
NCBI Protein Information	LIM domain kinase 2
UniProt Protein Name	LIM domain kinase 2
UniProt Gene Name	LIMK2
UniProt Synonym Gene Names	LIMK-2
UniProt Entry Name	LIMK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/limk-2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/319461>