

Rabbit MSLN Polyclonal Antibody | anti-MSLN antibody

MSLN Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA319507	Rabbit	Human, Rat, Mouse	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

MSLN; Polyclonal Antibody; MSLN Antibody; Mesothelin (CAK1 antigen; Pre-pro-megakaryocyte-potentiating factor)[Cleaved into: Megakaryocyte-potentiating factor (MPF); Mesothelin; cleaved form]; anti-MSLN antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	This antibody detects endogenous levels of Human MSLN
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-MSLN antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Immunogen	Synthesized peptide derived from human MSLN
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4 degree overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4 degree overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

ADDITIONAL INFORMATION

Related Product Information for anti-MSLN antibody	This gene encodes a preproprotein that is proteolytically processed to generate two protein products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiating factor functions as a cytokine that can stimulate colony formation of bone marrow megakaryocytes. Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016]
Product Categories/Family for anti-MSLN antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	53988378
NCBI GeneID	10232
NCBI Accession #	NP_005814.2
NCBI GenBank Nucleotide #	NM_005823.5
UniProt Accession #	Q13421; Q14859; Q4VQD5; Q96GR6; Q96KJ5; Q9BR17; Q9BTR2; Q9UCB2; Q9UK57; D3DU65
Molecular Weight	68,986 Da
NCBI Official Full Name	mesothelin isoform 1 preproprotein
NCBI Official Synonym Full Names	mesothelin
NCBI Official Symbol	MSLN
NCBI Official Synonym Symbols	MPF; SMRP
NCBI Protein Information	mesothelin; CAK1 antigen; megakaryocyte potentiating factor; soluble MPF mesothelin related protein; pre-pro-megakaryocyte-potentiating factor
UniProt Protein Name	Mesothelin
UniProt Gene Name	MSLN
UniProt Synonym Gene Names	MPF; MPF
UniProt Entry Name	MSLN_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/msln-antibody-immunohistochemistry-western-blot-elisa/319507>