

Mouse anti-Human Daxx Monoclonal Antibody | anti-Daxx antibody

Daxx Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA320032	Mouse	Human	ELISA, Flow Cytometry, Functional Assay, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

Daxx; Monoclonal Antibody; Daxx Monoclonal Antibody; Death domain-associated protein 6; DAXX; BING2; DAP6; hDaxx; ETS1-associated protein 1; EAP1; Fas death domain-associated protein; anti-Daxx antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Specificity	Daxx Monoclonal Antibody detects endogenous levels of Daxx protein.
Purity/Purification	Affinity purification
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-Daxx antibody	ELISA, FCM/FACS (Flow Cytometry), ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	Purified recombinant fragment of human Daxx expressed in E coli.
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of HeLa cells using Daxx Monoclonal Antibody (green) and negative control (purple).



IF (Immunofluorescence)

Confocal immunofluorescence analysis of PANC-1 cells using Daxx Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.



WB (Western Blot)
Western Blot analysis using Daxx Monoclonal Antibody against K562 cell lysate (1).

ADDITIONAL INFORMATION

Related Product Information for anti-Daxx antibody	This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript varian
Product Categories/Family for anti-Daxx antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	215422388
NCBI GeneID	1616
NCBI Accession #	NP_001135441.1
NCBI GenBank Nucleotide #	NM_001141969.1
UniProt Accession #	Q9UER7; O14747; O15141; O15208; Q5STK9; Q9BWI3; B4E1I3; F5H082
Molecular Weight	81,373 Da
NCBI Official Full Name	death domain-associated protein 6 isoform a
NCBI Official Synonym Full Names	death-domain associated protein
NCBI Official Symbol	DAXX
NCBI Official Synonym Symbols	DAP6; EAP1; BING2
NCBI Protein Information	death domain-associated protein 6; Fas-binding protein; CENP-C binding protein; ETS1-associated protein 1; death-associated protein 6; fas death domain-associated protein
UniProt Protein Name	Death domain-associated protein 6
UniProt Gene Name	DAXX
UniProt Synonym Gene Names	BING2; DAP6; hDaxx; EAP1
UniProt Entry Name	DAXX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/daxx-antibody-flow-cytometry-immunocytochemistry-immunofluorescence-western-blot-elisa/320032>