

Mouse SIRT1 Monoclonal Antibody | anti-SIRT1 antibody

SIRT1 Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA320225	Mouse	Human, Monkey	ELISA, Flow Cytometry, Functional Assay, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SIRT1; Monoclonal Antibody; SIRT1 Monoclonal Antibody; NAD-dependent deacetylase sirtuin-1; SIR2L1; NAD-dependent protein deacetylase sirtuin-1; hSIRT1; Regulatory protein SIR2 homolog 1; SIR2-like protein 1; hSIR2; anti-SIRT1 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Monkey
Clonality	Monoclonal
Specificity	SIRT1 Monoclonal Antibody detects endogenous levels of SIRT1 protein.
Purity/Purification	Affinity purification
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Applicable Applications for anti-SIRT1 antibody	ELISA, FCM/FACS (Flow Cytometry), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Purified recombinant fragment of human SIRT1 expressed in E coli.
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of K562 cells using SIRT1 Monoclonal Antibody (green) and negative control (purple).




IF (Immunofluorescence)

Immunofluorescence analysis of NTERA-2 cells using SIRT1 Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded lung cancer tissues (left) and kidney cancer tissues (right) with DAB staining using SIRT1 Monoclonal Antibody.



WB (Western Blot)

Western Blot analysis using SIRT1 Monoclonal Antibody against MCF-7 (1), Jurkat (2), HeLa (3), HEK293 (4) and A549 (5) cell lysate.

ADDITIONAL INFORMATION

Related Product Information for anti-SIRT1 antibody	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2008]
Product Categories/Family for anti-SIRT1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	215982798
NCBI GeneID	23411
NCBI Accession #	NP_001135970.1
NCBI GenBank Nucleotide #	NM_001142498.1
UniProt Accession #	Q96EB6; Q2XNF6; Q5JVQ0; Q9GZR9; Q9Y6F0
Molecular Weight	81,681 Da
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-1 isoform b
NCBI Official Synonym Full Names	sirtuin 1
NCBI Official Symbol	SIRT1
NCBI Official Synonym Symbols	SIR2L1
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-1; hSIR2; hSIRT1; SIR2alpha; sir2-like 1; sirtuin type 1; SIR2-like protein 1; regulatory protein SIR2 homolog 1; NAD-dependent deacetylase sirtuin-1
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-1
UniProt Gene Name	SIRT1
UniProt Synonym Gene Names	SIR2L1; hSIRT1; hSIR2; 75SirT1
UniProt Entry Name	SIR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

