

Mouse anti-Human OCT1 Monoclonal Antibody | anti-OCT1 antibody

Oct1 Monoclonal Antibody (7G1)

CATALOG # AAA320433	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	--

PRODUCT SYNONYMS

OCT1; Monoclonal Antibody; Oct1 Monoclonal Antibody (7G1); POU domain class 2 transcription factor; anti-OCT1 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Clone Number	7G1
Specificity	The antibody detects endogenous Oct1 protein.
Purity/Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Form/Format	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Applicable Applications for anti-OCT1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthetic Peptide of 42644
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of Human-breast-cancer tissue. 1.Oct1 Monoclonal Antibody(7G1)(red) was diluted at 1:200(4 degree C.overnight). 2. Cy3 labeled Secondary antibody was diluted at 1:300(room temperature. 50min).3. Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human-breast-cancer tissue. 1.Oct1 Monoclonal Antibody(7G1) was diluted at 1:200(4 degree C.overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98 degree C.20min). 3.Secondary antibody was diluted at 1:200(room temperature. 30min). Negative control was used by secondary antibody only.



WB (Western Blot)

Western blot analysis of 1) Hela. 2) Jurkat. 3) HepG2. diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003.Inventbiotech.MN.USA).

ADDITIONAL INFORMATION

Related Product Information for anti-OCT1 antibody	The OCT1 transcription factor was among the first identified members of the POU transcription factor family (summarized by Sturm et al., 1993 [PubMed 8314572]). Members of this family contain the POU domain, a 160-amino acid region necessary for DNA binding to the octameric sequence ATGCAAAT.[supplied by OMIM, Jul 2010]
Product Categories/Family for anti-OCT1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	311771667
NCBI GeneID	5451
NCBI Accession #	NP_001185712.1
NCBI GenBank Nucleotide #	NM_001198783.1
UniProt Accession #	P14859; P09086
Molecular Weight	76,472 Da
NCBI Official Full Name	POU domain, class 2, transcription factor 1 isoform 2
NCBI Official Synonym Full Names	POU class 2 homeobox 1
NCBI Official Symbol	POU2F1
NCBI Official Synonym Symbols	OCT1; OTF1; oct-1B
NCBI Protein Information	POU domain, class 2, transcription factor 1; NF-A1; OTF-1; oct-1; octamer-binding protein 1; Octamer-binding transcription factor-1; octamer-binding transcription factor 1
UniProt Protein Name	POU domain, class 2, transcription factor 1
UniProt Gene Name	POU2F1
UniProt Synonym Gene Names	OCT1; OTF1; Oct-1; OTF-1
UniProt Entry Name	PO2F1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.