

Rabbit hnRPD Polyclonal Antibody | anti-hnRPD antibody

Phospho-hnRPD(Ser83) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA320733	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

hnRPD; Polyclonal Antibody; Phospho-hnRPD(Ser83) Antibody; ARE binding protein AUF1 type A; AU-rich element RNA-binding protein 1; AUF; AUF1; AUF1A; Heterogeneous nuclear ribonucleoprotein D0; hnRNP D; hnRNP D0; Hnnpd; hnRNPD0; HNRPD; HNRPD_HUMAN; P37; anti-hnRPD antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-hnRPD(Ser83) antibody detects endogenous levels of hnRPD only when phosphorylated at Sersine 83
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	287
Applicable Applications for anti-hnRPD antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human hnRPD around the phosphorylation site of Sersine 83
Subcellular Location	Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Component of ribonucleosomes.
Conjugation	Unconjugated
Epitope	Phospho Ser83
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)

AAA320733 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



IHC (Immunohistochemistry)

AAA320733 at 1/200 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IF (Immunofluorescence)

AAA320733 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



WB (Western Blot)

Western blot analysis on HuvEc cell lysate using Phospho-hnRPD(Ser83) Antibody, The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-hnRPD antibody

Description: This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are nucleic acid binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It localizes to both the nucleus and the cytoplasm. This protein is implicated in the regulation of mRNA stability. Alternative splicing of this gene results in four transcript variants.

Function: Binds with high affinity to RNA molecules that contain AU-rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Also binds to double- and single-stranded DNA sequences in a specific manner and functions as a transcription factor. Each of the RNA-binding domains specifically can bind solely to a single-stranded non-monotonous 5'-UUAG-3' sequence and also weaker to the single-stranded 5'-TTAGGG-3' telomeric DNA repeat. Binds RNA oligonucleotides with 5'-UUAGGG-3' repeats more tightly than the telomeric single-stranded DNA 5'-TTAGGG-3' repeats. Binding of RRM1 to DNA inhibits the formation of DNA quadruplex structure which may play a role in telomere elongation. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. May play a role in the regulation of the rhythmic expression of circadian clock core genes. Directly binds to the 3'UTR of CRY1 mRNA and induces CRY1 rhythmic translation. May also be involved in the regulation of PER2 translation.

Subunit Structure: Identified in a IGF2BP1-dependent mRNP granule complex containing untranslated mRNAs. Part of a complex associated with the FOS mCRD domain and consisting of PABPC1, PAIP1, CSDE1/UNR and SYNCRIP. Interacts with IGF2BP2. Interacts with GTPBP1. Interacts with EIF4G1; the interaction requires RNA. Interacts with EIF3B and RPS3.

Post-translational Modifications: Arg-345 is dimethylated, probably to asymmetric dimethylarginine. Methylated by PRMT1, in an insulin-dependent manner. The PRMT1-mediated methylation regulates tyrosine phosphorylation (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	51477708
NCBI GeneID	3184
NCBI Accession #	NP_001003810.1
NCBI GenBank Nucleotide #	NM_001003810.1
UniProt Accession #	Q14103 ; P07029 ; Q01858 ; Q14100 ; Q14101 ; Q14102 ; Q4W5A1 ; Q9UCE8 ; Q9UCE9 ; A8K9J2
Molecular Weight	Observed: 38 kDa Predicted: 39 kDa
NCBI Official Full Name	heterogeneous nuclear ribonucleoprotein D0 isoform d
NCBI Official Synonym Full Names	heterogeneous nuclear ribonucleoprotein D
NCBI Official Symbol	HNRNPD
NCBI Official Synonym Symbols	P37; AUF1; AUF1A; HNRPD; hnRNPD0
NCBI Protein Information	heterogeneous nuclear ribonucleoprotein D0

UniProt Protein Name	Heterogeneous nuclear ribonucleoprotein D0
UniProt Gene Name	HNRNPD
UniProt Synonym Gene Names	AUF1; HNRPD; hnRNP D0

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hnrpd-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/320733>