

Rabbit GLP1 Polyclonal Antibody | anti-GLP1 antibody

GLP1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA320811	Rabbit	Human, Mouse, Rat, Monkey	ELISA, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS

GLP1; Polyclonal Antibody; GLP1 Antibody; GCG; GLP-1; GLP-1(7-36); GLP-1(7-37); GLP-2; GLUC_HUMAN; Glucagon-like peptide 2; GRPP; Incretin hormone; OXM; OXY; anti-GLP1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	GLP1 Antibody detects endogenous levels of total GLP1.
Purity/Purification	Peptide affinity purification
Form/Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-GLP1 antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human GLP1, corresponding to a region within the internal amino acids.
Preparation and Storage	Store at -20 °C. Stable for 12 months from date of receipt.

IMAGES




IHC (Immunohistochemistry)

AAA320811 at 1/100 staining Human pancreatic cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



WB (Western Blot)

Western blot analysis on COS7 cell lysate using Glucagon Antibody, The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)

AAA320811 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/glp1-antibody-immunohistochemistry-immunocytochemistry-immunofluorescence-western-blot-elisa/320811>