

Rabbit GK Polyclonal Antibody | anti-GK antibody

GK Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA321249	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS

GK; Polyclonal Antibody; GK Antibody; ATP glycerol 3 phosphotransferase; ATP:glycerol 3 phosphotransferase; ATP:glycerol 3-phosphotransferase; D930012N15Rik; GK1; GKD; GLPK_HUMAN; Glycerokinase; Glycerol kinase; OTTHUMP00000023108; OTTHUMP00000023109; OTTHUMP00000215321; anti-GK antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	GK antibody detects endogenous levels of GK
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	524
Applicable Applications for anti-GK antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human GK
Subcellular Location	Mitochondrion Outer Membrane. Cytoplasm. In sperm and fetal tissues, the majority of the enzyme is bound to mitochondria, but in adult tissues, such as liver found in the cytoplasm.
Tissue Specificity	Highly expressed in the liver, kidney and testis. Isoform 2 and isoform 3 are expressed specifically in testis and fetal liver, but not in the adult liver.
Predicted Cross Reactivity	Pig, Bovine, Sheep, Rabbit, Dog
Similarity	Pig (100%), Bovine (100%), Sheep (100%), Rabbit (100%), Dog (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western blot analysis on 293 cell lysate using GK Antibody. The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)

AAA321249 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-GK antibody	Description: Glycerol kinase (EC 2.7.1.30) catalyzes the phosphorylation of glycerol by ATP, yielding ADP and glycerol-3-phosphate.[supplied by OMIM] Function: Key enzyme in the regulation of glycerol uptake and metabolism. Similarity: Belongs to the FGGY kinase family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504007
NCBI GeneID	2710
NCBI Accession #	NP_000158.1
NCBI GenBank Nucleotide #	NM_000167.5
UniProt Accession #	P32189; Q6IQ27; Q8IVR5; Q9UMP0; Q9UMP1; A6NJP5; B2R833
Molecular Weight	Observed: 57 kDa Predicted: 62 kDa
NCBI Official Full Name	glycerol kinase isoform b
NCBI Official Synonym Full Names	glycerol kinase
NCBI Official Symbol	GK
NCBI Official Synonym Symbols	GK1; GKD
NCBI Protein Information	glycerol kinase
UniProt Protein Name	Glycerol kinase
UniProt Gene Name	GK
UniProt Synonym Gene Names	GK; Glycerokinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gk-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/321249>