

Rabbit FOS Polyclonal Antibody | anti-FOS antibody

Phospho-FOS (Ser32) Antibody

CATALOG # AAA321287	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

FOS; Polyclonal Antibody; Phospho-FOS (Ser32) Antibody; Activator protein 1; AP 1; C FOS; Cellular oncogene c fos; Cellular oncogene fos; FBJ murine osteosarcoma viral (v fos) oncogene homolog (oncogene FOS); FBJ murine osteosarcoma viral oncogene homolog; FBJ murine osteosarcoma viral v fos oncogene homolog; FBJ Osteosarcoma Virus; FOS protein; FOS_HUMAN; G0 G1 switch regulatory protein 7; G0/G1 switch regulatory protein 7; G0S7; Oncogene FOS; p55; proto oncogene c Fos; Proto oncogene protein c fos; Proto-oncogene c-Fos; v fos FBJ murine osteosarcoma viral oncogene homolog; anti-FOS antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-FOS (Ser32) antibody detects endogenous levels of FOS.
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	380
Applicable Applications for anti-FOS antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human FOS around the phosphorylation site of Ser32.
Subcellular Location	Nucleus.
Tissue Specificity	Expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, they undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the protein becomes nuclear.
Predicted Cross Reactivity	Pig, Bovine, Horse, Rabbit, Dog, Chicken
Similarity	Pig (100%), Bovine (100%), Horse (100%), Rabbit (100%), Dog (100%), Chicken (83%)
Conjugation	Unconjugated
Epitope	Phospho Ser32
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
AAA321287 at 1/100 staining Mouse lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of extracts of HeLa serum-starved treated with TPA, using Phospho-FOS (Ser32) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-FOS antibody	Function: Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, FOS and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. On TGF-beta activation, forms a multimeric SMAD3/SMAD4/JUN/FOS complex at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation. In growing cells, activates phospholipid synthesis, possibly by activating CDS1 and PI4K2A. This activity requires Tyr-dephosphorylation and association with the endoplasmic reticulum. Subunit Structure: Heterodimer; with JUN (By similarity). Interacts with MAFB (By similarity). Component of the SMAD3/SMAD4/JUN/FOS complex required for synergistic TGF-beta-mediated transcription at the AP1 promoter site. Interacts with SMAD3; the interaction is weak even on TGF-beta activation. Interacts with MAFB. Interacts with DSIP1; this interaction inhibits the binding of active AP1 to its target DNA. Interacts with CDS1 and PI4K2A (By similarity). Post-translational Modifications: Phosphorylated in the C-terminal upon stimulation by nerve growth factor (NGF) and epidermal growth factor (EGF). Phosphorylated, in vitro, by MAPK and RSK1. Phosphorylation on both Ser-362 and Ser-374 by MAPK1/2 and RSK1/2 leads to protein stabilization with phosphorylation on Ser-374 being the major site for protein stabilization on NGF stimulation. Phosphorylation on Ser-362 and Ser-374 primes further phosphorylations on Thr-325 and Thr-331 through promoting docking of MAPK to the DEF domain. Phosphorylation on Thr-232, induced by HA-RAS, activates the transcriptional activity and antagonizes sumoylation. Phosphorylation on Ser-362 by RSK2 in osteoblasts contributes to osteoblast transformation (By similarity). Constitutively sumoylated with SUMO1, SUMO2 and SUMO3. Desumoylated by SENP2. Sumoylation requires heterodimerization with JUN and is enhanced by mitogen stimulation. Sumoylation inhibits the AP-1 transcriptional activity and is, itself, inhibited by Ras-activated phosphorylation on Thr-232. In quiescent cells, the small amount of FOS present is phosphorylated at Tyr-10 and Tyr-30 by SRC. This Tyr-phosphorylated form is cytosolic. In growing cells, dephosphorylated by PTPN2. Dephosphorylation leads to the association with endoplasmic reticulum membranes and activation of phospholipid synthesis. Similarity: Belongs to the bZIP family. Fos subfamily.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885241
NCBI GeneID	2353
NCBI Accession #	NP_005243.1
NCBI GenBank Nucleotide #	NM_005252.3
UniProt Accession #	P01100; P18849; A8K4E2; B4DQ65
Molecular Weight	Observed: 62 kDa Predicted: 41 kDa

NCBI Official Full Name	proto-oncogene c-Fos
NCBI Official Synonym Full Names	Fos proto-oncogene, AP-1 transcription factor subunit
NCBI Official Symbol	FOS
NCBI Official Synonym Symbols	p55; AP-1; C-FOS
NCBI Protein Information	proto-oncogene c-Fos
UniProt Protein Name	Proto-oncogene c-Fos
UniProt Gene Name	FOS
UniProt Synonym Gene Names	G0S7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fos-antibody-immunohistochemistry-western-blot-elisa/321287>