

Rabbit DDR1 Polyclonal Antibody | anti-DDR1 antibody

Phospho-DDR1(Tyr513) Antibody

CATALOG # AAA321296	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---

PRODUCT SYNONYMS


DDR1; Polyclonal Antibody; Phospho-DDR1(Tyr513) Antibody; CAK; CD 167; CD167; CD167 antigen-like family member A; CD167a; Cell adhesion kinase; DDR 1; DDR; DDR1_HUMAN; Discoidin domain receptor; Discoidin domain receptor tyrosine kinase 1; Discoidin receptor tyrosine kinase; Discoidin receptor tyrosine kinase isoform a; EDDR 1; EDDR1; Epithelial discoidin domain receptor 1; Epithelial discoidin domain-containing receptor 1; Epithelial specific receptor kinase; HGK2; Mammarian carcinoma kinase 10; Mammary carcinoma kinase 10; MCK-10; MCK10; NEP; Neuroepithelial tyrosine kinase; Neurotrophic tyrosine kinase receptor type 4; NTRK 4; NTRK4; OTTHUMP00000029343; OTTHUMP00000029344; OTTHUMP00000029345; OTTHUMP00000029346; OTTHUMP00000029347; OTTHUMP00000164863; OTTHUMP00000164867; OTTHUMP00000222080; Protein-tyrosine kinase 3A; Protein-tyrosine kinase RTK-6; PTK 3; PTK 3A protein tyrosine kinase 3A; PTK3; PTK3A; Receptor tyrosine kinase NEP; RTK 6; RTK6; TRK E; TRKE; Tyrosine kinase DDR; Tyrosine kinase receptor E; Tyrosine-protein kinase CAK; anti-DDR1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-DDR1(Tyr513) antibody detects endogenous levels of DDR1.
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	508
Applicable Applications for anti-DDR1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human DDR1 around the phosphorylation site of Tyr513.
Subcellular Location	Secreted and Membrane.
Tissue Specificity	Detected in T-47D, MDA-MB-175 and HBL-100 breast carcinoma cells, A-431 epidermoid carcinoma cells, SW48 and SNU-C2B colon carcinoma cells and Hs 294T melanoma cells (at protein level). Expressed at low levels in most adult tissues and is highest in the brain, lung, placenta and kidney. Lower levels of expression are detected in melanocytes, heart, liver, skeletal muscle and pancreas. Abundant in breast carcinoma cell lines. In the colonic mucosa, expressed in epithelia but not in the connective tissue of the lamina propria. In the thyroid gland, expressed in the epithelium of the thyroid follicles. In pancreas, expressed in the islets of Langerhans cells, but not in the surrounding epithelial cells of the exocrine pancreas. In kidney, expressed in the epithelia of the distal tubules. Not expressed in connective tissue, endothelial cells, adipose tissue, muscle cells or cells of hematopoietic origin.
Predicted Cross Reactivity	Horse, Sheep, Rabbit
Similarity	Horse (100%), Sheep (100%), Rabbit (100%)
Conjugation	Unconjugated

Epitope	Phospho Tyr513
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohiostchemistry)
AAA321296 at 1/100 staining Human breast cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of extracts of T47-D cells treated with collagen, using Phospho-DDR1(Tyr513) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-DDR1 antibody	Function: Tyrosine kinase that functions as cell surface receptor for fibrillar collagen and regulates cell attachment to the extracellular matrix, remodeling of the extracellular matrix, cell migration, differentiation, survival and cell proliferation. Collagen binding triggers a signaling pathway that involves SRC and leads to the activation of MAP kinases. Regulates remodeling of the extracellular matrix by up-regulation of the matrix metalloproteinases MMP2, MMP7 and MMP9, and thereby facilitates cell migration and wound healing. Required for normal blastocyst implantation during pregnancy, for normal mammary gland differentiation and normal lactation. Required for normal ear morphology and normal hearing (By similarity). Promotes smooth muscle cell migration, and thereby contributes to arterial wound healing. Also plays a role in tumor cell invasion. Phosphorylates PTPN11. Subunit Structure: Homodimer. Interacts (via PPxY motif) with WWC1 (via WW domains) in a collagen-regulated manner. Forms a tripartite complex with WWC1 and PRKCZ, but predominantly in the absence of collagen. Interacts (tyrosine phosphorylated) with SHC1. Interacts with SRC. Interacts with MYH9. Interacts with CDH1. Interacts with PTPN11. Interacts with NCK2. Post-translational Modifications: Autophosphorylated in response to fibrillar collagen binding. Glycosylation of Asn-211, but apparently not of Asn-260, Asn-371, or Asn-394, prevents autophosphorylation from occurring in the absence of collagen. Similarity: The Gly/Pro-rich domains may be required for an unusual geometry of interaction with ligand or substrates. Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	321400059
NCBI GeneID	780
NCBI Accession #	NP_001189450.1
NCBI GenBank Nucleotide #	NM_001202521.1
UniProt Accession #	Q08345; Q14196; Q16562; Q2L6H3; Q4LE50; Q5ST11; Q5ST12; Q6NSK4; Q9UD35; B5A975; B5A976; B7Z2K0
Molecular Weight	Observed: 125 kDa Predicted: 102 kDa
NCBI Official Full Name	epithelial discoidin domain-containing receptor 1 isoform 4
NCBI Official Synonym Full Names	discoidin domain receptor tyrosine kinase 1
NCBI Official Symbol	DDR1

NCBI Official Synonym Symbols	CAK; DDR; NEP; HGK2; PTK3; RTK6; TRKE; CD167; EDDR1; MCK10; NTRK4; PTK3A
NCBI Protein Information	epithelial discoidin domain-containing receptor 1
UniProt Protein Name	Epithelial discoidin domain-containing receptor 1
UniProt Gene Name	DDR1
UniProt Synonym Gene Names	CAK; EDDR1; NEP; NTRK4; PTK3A; RTK6; TRKE; Epithelial discoidin domain receptor 1; MCK-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ddr1-antibody-immunohistochemistry-western-blot-elisa/321296>