

Rabbit anti-Human IRF-3 Polyclonal Antibody | anti-IRF-3 antibody

Phospho-IRF-3 (Ser396) Antibody


CATALOG # AAA321337	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---

PRODUCT SYNONYMS


IRF-3; Polyclonal Antibody; Phospho-IRF-3 (Ser396) Antibody; IIAE7; Interferon regulatory factor 3; IRF 3; IRF3; IRF3_HUMAN; MGC94729; anti-IRF-3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-IRF-3 (Ser396) antibody detects endogenous levels of IRF-3.
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	452
Applicable Applications for anti-IRF-3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human IRF-3 around the phosphorylation site of Ser396.
Subcellular Location	Cytoplasm. Nucleus. Shuttles between cytoplasmic and nuclear compartments, with export being the prevailing effect. When activated, IRF3 interaction with CREBBP prevents its export to the cytoplasm.
Tissue Specificity	Expressed constitutively in a variety of tissues.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Dog
Similarity	Pig (100%), Bovine (91%), Horse (100%), Sheep (91%), Dog (82%)
Conjugation	Unconjugated
Epitope	Phospho Ser396
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



IHC (Immunohistochemistry)
AAA321337 at 1/100 staining Human lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of extracts of HT-29 cells transfected with Poly (I:C), using Phospho-IRF-3 (Ser396) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-IRF-3 antibody	Function: Key transcriptional regulator of type I interferon (IFN)-dependent immune responses which plays a critical role in the innate immune response against DNA and RNA viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, is phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAFI), a complex which activates the transcription of the type I IFN and ISG genes. Can activate distinct gene expression programs in macrophages and can induce significant apoptosis in primary macrophages. Subunit Structure: Monomer. Homodimer; phosphorylation-induced. Heterodimer with IRF7. Interacts with CREBBP. May interact with MAVS. Interacts with IKBKE and TBK1. Interacts with TICAM1 and TICAM2. Interacts with RBCK1. Interacts with HERC5. Interacts with DDX3X (phosphorylated at 'Ser-102'); the interaction allows the phosphorylation and activation of IRF3 by IKBKE. Interacts with TRIM21 and ULK1, in the presence of TRIM21; this interaction leads to IRF3 degradation by autophagy (PubMed:26347139). Interacts with RIOK3; RIOK3 probably mediates the interaction of TBK1 with IRF3. Post-translational Modifications: Constitutively phosphorylated on many Ser/Thr residues. C-terminal serine/threonine cluster is phosphorylated in response of induction by IKBKE and TBK1. Phosphorylated at Ser-396 by IKBKE upon ssRNA viral infection. Ser-385 and Ser-386 may be specifically phosphorylated in response to induction. Phosphorylation at Ser-386 by TBK1 results in oligomerization. An alternate model propose that the five serine/threonine residues between 396 and 405 are phosphorylated in response to a viral infection. (Microbial infection) Phosphorylation and subsequent activation of IRF3 is inhibited by vaccinia virus protein E3. Ubiquitinated; ubiquitination involves RBCK1 leading to proteasomal degradation. Polyubiquitinated; ubiquitination involves TRIM21 leading to proteasomal degradation. ISGylated by HERC5 resulting in sustained IRF3 activation and in the inhibition of IRF3 ubiquitination by disrupting PIN1 binding. The phosphorylation state of IRF3 does not alter ISGylation. Similarity: Belongs to the IRF family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	308199446
NCBI GeneID	3661
NCBI Accession #	NP_001184051.1
NCBI GenBank Nucleotide #	NM_001197122.1
UniProt Accession #	Q14653; Q5FBY1; Q5FBY2; Q5FBY4; Q7Z5G6; A8K7L2; B2RAZ3

Molecular Weight	Observed: 45-55 kDa Predicted: 48 kDa
NCBI Official Full Name	interferon regulatory factor 3 isoform 2
NCBI Official Synonym Full Names	interferon regulatory factor 3
NCBI Official Symbol	IRF3
NCBI Official Synonym Symbols	IIAE7
NCBI Protein Information	interferon regulatory factor 3
UniProt Protein Name	Interferon regulatory factor 3
UniProt Gene Name	IRF3
UniProt Synonym Gene Names	IRF-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/irf-3-antibody-immunohistochemistry-western-blot-elisa/321337>