

Rabbit IRF3 Polyclonal Antibody | anti-IRF3 antibody

Phospho-IRF3(Ser396) Antibody

CATALOG # AAA321418	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

IRF3; Polyclonal Antibody; Phospho-IRF3(Ser396) Antibody; IIAE7; Interferon regulatory factor 3; IRF 3; IRF-3; IRF3_HUMAN; MGC94729; anti-IRF3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Sequence Length	452
Applicable Applications for anti-IRF3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human IRF3 around the phosphorylation site of Ser396.
Preparation and Storage	Store at -20°C. Stable for 12 months from date of receipt.

IMAGES




WB (Western Blot)

Western blot analysis of extracts from HT29 cells plpC-transfected, using Phospho-IRF3(Ser396) Antibody.



IHC (Immunohistochemistry)

AAA321418 at 1/100 staining Mouse thymus tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IF (Immunofluorescence)

AAA321418 staining HepG2 cells(30min of 4uM Forskolin treatment) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AAA321418) and mouse anti-beta tubulin Ab for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.The nuclear counter stain is DAPI(blue).

ADDITIONAL INFORMATION

Related Product Information for anti-IRF3 antibody

Function: Key transcriptional regulator of type I interferon (IFN)-dependent immune responses which plays a critical role in the innate immune response against DNA and RNA viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, is phosphorylated by IKBKE and TBK1 kinases.

NCBI AND UNIPROT INFORMATION

NCBI GI #	308199446
NCBI GeneID	3661
NCBI Accession #	NP_001184051.1
NCBI GenBank Nucleotide #	NM_001197122.1
UniProt Accession #	Q14653; Q5FBY1; Q5FBY2; Q5FBY4; Q7Z5G6; A8K7L2; B2RAZ3
Molecular Weight	45-55 kDaa
NCBI Official Full Name	interferon regulatory factor 3 isoform 2
NCBI Official Synonym Full Names	interferon regulatory factor 3
NCBI Official Symbol	IRF3
NCBI Official Synonym Symbols	IIAE7
NCBI Protein Information	interferon regulatory factor 3
UniProt Protein Name	Interferon regulatory factor 3
UniProt Gene Name	IRF3
UniProt Synonym Gene Names	IRF-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.